

# METHODS OF MATHEMATICAL PHYSICS

## Infinite series; Residues; Analytic Continuation

## Tutorial Sheet 1

**K:** key question – explores core material

**R:** review question – an invitation to consolidate

**C:** challenge question – going beyond the basic framework of the course

**S:** standard question – general fitness training!

### 1.1 ‘Telescoping’ an infinite series [S]

$$\text{Consider } S = \sum_{n=1}^{\infty} \frac{n}{(n+1)!}$$

By writing  $n = (n+1) - 1$  in the numerator of the summand, show that a large cancellation occurs between the resulting two series. Hence evaluate  $S$ .

### 1.2 Comparison of a series with an integral [S]

By comparison with an integral, show that the series

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^s}$$

converges for  $s > 1$ .

What does the ratio test yield for this series?

### 1.3 Bernoulli Numbers [C]

Consider the infinite series

$$\frac{x}{e^x - 1} = B_0 + B_1x + \frac{B_2}{2!}x^2 \cdots + \frac{B_n}{n!}x^n \cdots$$

By expanding the lhs and equating powers of  $x$  determine that

$$\begin{aligned} 1 &= B_0 \\ 0 &= \frac{B_0}{2!} + \frac{B_1}{1!} \\ 0 &= \frac{B_0}{3!} + \frac{B_1}{2! 1!} + \frac{B_2}{1! 2!} \end{aligned}$$

Show that for  $n > 1$  one can write the relations as

$$(B+1)^n - B^n = 0 \quad \text{where } B^s \rightarrow B_s$$

.

### 1.4 Asymptotic expansion of error function [K]

The error function is defined as

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x dt e^{-t^2}$$

Show by considering the complementary error function  $\text{erfc}(x) = 1 - \text{erf}(x)$  and integrating by parts that

$$\text{erf}(x) \sim 1 - \frac{1}{\sqrt{\pi}} \frac{e^{-x^2}}{x} \sum_{n=0}^{\infty} \frac{(-1)^n (2n-1)!!}{x^{2n} 2^n}$$

where  $(2n-1)!! = (2n-1)(2n-3)\cdots(3)(1)$

Explain why this is an asymptotic expansion of the error function.

### 1.5 Analytic Continuation [K]

Suppose we are given the power series expansion

$$f(z) = - \sum_{n=1}^{\infty} \frac{z^n}{n}$$

which converges for  $|z| < 1$ .

Sum this series and derive a power series expansion for the resulting function about  $z = -1/2$ . What is the radius of convergence of this series?

Repeat for expansions about  $z = 1/2$  and  $z = 3/4$ .

Discuss the complication that arises when analytically continuing to the point  $z = 2$ .

### 1.6 Integrating via the residue method [S]

Compute the following real or complex integrals by using the residue method:

(i)

$$\int_{-\infty}^{+\infty} \frac{e^{iax}}{a^2 + x^2} dx,$$

where  $a$  is a real number.

(ii)

$$\oint_{\gamma} \frac{dz}{z^{n-m+1}},$$

where  $n$  and  $m$  are integers, and  $\gamma$  is a circuit enclosing the origin once in an anti-clockwise direction.

(iii)

$$\int_0^{2\pi} \frac{d\theta}{5 - 4 \cos(\theta)},$$

To solve this, you should use the fact that  $\cos(\theta) = \frac{e^{i\theta} + e^{-i\theta}}{2}$ .

### 1.7 Bernoulli numbers from a contour integration [C]

(i) Show that generalising  $x$  to complex variable  $z$  (analytic continuation) in question 1.3 implies that

$$B_n = \frac{n!}{2\pi i} \oint_C \frac{z}{e^z - 1} \frac{dz}{z^{n+1}}$$

where the contour  $C$  encircles the origin in an anticlockwise fashion with  $|z| < 2\pi$  to avoid the poles at  $\pm 2\pi i$ .

(ii) Locate and classify all the singularities of the integrand.

(iii) By the residue theorem  $B_n = 2\pi i \times$  residue from origin. By deforming the contour show that

$$B_n = -2\pi i \times \text{sum of all other residues}$$

Hint: you will need to consider a large circle of radius  $R \rightarrow \infty$  encircling the origin in a *clockwise* fashion

(iv) Hence show that

$$\begin{aligned} B_n &= 0 \quad \text{for } n \text{ odd} \\ B_n &= -2 \frac{(-1)^{n/2} n!}{(2\pi)^n} \zeta(n) \quad \text{for } n \text{ even} \end{aligned}$$

where  $\zeta(s) = \sum_{m=1}^{\infty} m^{-s}$ . By calculating explicitly the Bernoulli numbers in 1.3, evaluate  $\zeta(2)$  and  $\zeta(4)$