

Symmetries of Classical Mechanics (SoCM) – Example sheet VII: 11.11.11

Semester 1 2011-2012

1. Evaluate the integrals

(a)

$$I = \int_{\text{all space}} e^{-a_{ij}x_i x_j} dV ,$$

where a_{ij} are the elements of a real symmetric positive definite matrix.
[It may be assumed without proof that $\int_{-\infty}^{\infty} d\xi e^{-a\xi^2} = \sqrt{\frac{\pi}{a}}$ for $\text{Re}(a) > 0$.]
[Hint: rotate to the principal axis basis.]

(b)

$$A_{ij} = \int_{\text{sphere } r \leq a} x_i x_j dV .$$

[Hint: first argue that A_{ij} is a rank two isotropic tensor and then, for example, consider the sum over $i = j$.]

(c)

$$A_{ijkl} = \int_{\text{sphere } r \leq a} x_i x_j x_k x_l dV .$$

[Hint: first argue that A_{ijkl} is a rank four isotropic tensor and then proceed as in the previous part.]

2. A piece of material is subjected to a homogeneous stress such that the forces per unit area across faces with (orthogonal) unit normals \underline{e}_1 , \underline{e}_2 and \underline{e}_3 are $(5p\underline{e}_1 + 2p\underline{e}_3)$, $p\underline{e}_2$ and $(2p\underline{e}_1 + 2p\underline{e}_3)$ respectively. Construct the (homogeneous) stress tensor within the material and calculate the force acting across the element of area δS whose normal is in the direction $(\underline{e}_1 + 2\underline{e}_2 + 2\underline{e}_3)$. Find the principal axes of stress and the corresponding principal stresses.

3. If the stress-strain relation is given by

$$P_{ij} = \lambda \text{Tr} E \delta_{ij} + 2\mu E_{ij} ,$$

then by considering a simple tension/unit area S in an axially symmetric rod placed in the x_1 direction, show that this may alternatively be written as

$$P_{ij} = \frac{Y}{(1 + \sigma)} \left[\frac{\sigma}{(1 - 2\sigma)} \text{Tr} E \delta_{ij} + E_{ij} \right] ,$$

where Y is Young's modulus, defined by $Y = S/E_{11}$ and σ is Poisson's ratio, defined by $\sigma = -E_{22}/E_{11} \equiv -E_{33}/E_{11}$.

[Hint: look at your notes.]

4. The point (x_1, x_2, x_3) in an isotropic, homogeneous material is displaced by a small amount (u_1, u_2, u_3) given by

$$u_1 = -\frac{\sigma}{Y}\alpha x_1 + \frac{(1+\sigma)}{Y}\beta x_2, \quad u_2 = \frac{(1+\sigma)}{Y}\beta x_1 - \frac{\sigma}{Y}\alpha x_2, \quad u_3 = \frac{1}{Y}\alpha x_3,$$

where Y and σ are Young's modulus and Poisson's ratio respectively, and α and β are constants. Evaluate the strain tensor, and hence find the nature and magnitude of the stresses that are causing this deformation.

5. The point (x_1, x_2, x_3) in an isotropic, homogeneous medium undergoes a small displacement (u_1, u_2, u_3) given by

$$u_1 = \alpha x_1, \quad u_2 = -3d x_2 - d x_3, \quad u_3 = -d x_2 - 3d x_3,$$

where α and d are constants. Write down the strain tensor and find the principal strains β and γ ($|\beta| < |\gamma|$) in the \underline{e}_2 - \underline{e}_3 plane.

Show that this strain may be maintained by a stress P_1 along the \underline{e}_1 axis, a stress P_2 along the principal axis of strain which has principal strain β , and zero stress along the third principal axis, provided that $d = \sigma\alpha/(4-2\sigma)$, where σ is Poisson's ratio for the medium.

6. A bar of homogeneous, isotropic material has a rectangular cross-section. It is subjected to a uniform, normal pressure P over its four sides, and the end faces are subjected to a tension αP per unit area acting parallel to the axis of the bar. Find the ratio of the lateral strain to the longitudinal strain in terms of α and Poisson's ratio, σ , for the material of the bar. Show that this ratio is independent of σ for $\alpha = 2$ or $\alpha = -1$ and state the corresponding values of the ratio.
7. (a) If \underline{u} is a vector field, show that $\underline{\nabla} \cdot \underline{u}$ is a scalar field.
 (b) If ϕ is a scalar field, show that the quantity

$$T_{ij} = \frac{\partial^2 \phi}{\partial x_i \partial x_j},$$

is a second rank tensor.