

Physics Prospects at the HL-LHC



LHC Run 1 (& 2)

IN



proton-proton collisions at ATLAS and CMS

- ▶ 2010 $\sqrt{s}=7$ TeV, 44 pb^{-1}
- ▶ 2011 $\sqrt{s}=7$ TeV, 6 fb^{-1}
- ▶ 2012 $\sqrt{s}=8$ TeV, 23 fb^{-1}
- ▶ Run 2: 2015 $\sqrt{s}=13$ TeV, 4 fb^{-1}

OUT

Physics results!

- ▶ Nearly 1000 submitted papers on Run 1 collision data
- ▶ 9 papers on Run 2 data



(generated 2016-01-29 09:22 including fill 4569)

The Nobel Prize in Physics 2013

François Englert and Peter W. Higgs



"for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN's Large Hadron Collider"

Run 1 Higgs Boson Results

All observations from the LHC consistent with a Standard Model Higgs boson with $m_H \sim 125$ GeV.

➔ m_H measured in ZZ and $\gamma\gamma$ final states consistent with 125 GeV.

➔ It decays like a SM Higgs boson



➔ It's produced like a SM Higgs boson

[arXiv:1412.8662](https://arxiv.org/abs/1412.8662)

[Phys. Rev. D. 90, 052004 \(2014\)](https://arxiv.org/abs/1412.8662)

[ATLAS-CONF-044](https://arxiv.org/abs/1412.8662)


[Eur. Phys. J. C 74 \(2014\) 3076](https://arxiv.org/abs/1412.8662)




But not only ...

Standard Model Production Cross Section Measurements

Status: Nov 2015



95% CL Limits on Masses of Exotic Phenomena in TeV



ATLAS SUSY Searches* - 95% CL Lower Limits

Status: July 2015

ATLAS Preliminary

$\sqrt{s} = 7, 8 \text{ TeV}$

Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L} d\sqrt{s} [\text{fb}^{-1}]$	Mass limit	$\sqrt{s} = 7 \text{ TeV}$	$\sqrt{s} = 8 \text{ TeV}$	Reference	
Inclusive Searches	MSUGRA/CMSSM	0-3 $e, \mu/1-2 \tau$	2-10 jets/3 b	Yes	20.3	\tilde{q}, \tilde{g}	1.8 TeV	$m(\tilde{q})=m(\tilde{g})$	1507.05525
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{q}	850 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(1^{\text{st}} \text{ gen. } \tilde{q})=m(2^{\text{nd}} \text{ gen. } \tilde{q})$	1405.7875
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ (compressed)	mono-jet	1-3 jets	Yes	20.3	\tilde{q}	100-440 GeV	$m(\tilde{q})-m(\tilde{\chi}_1^0)<10 \text{ GeV}$	1507.05525
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ (off-Z)	2 jets	Yes	20.3	\tilde{q}	780 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	1503.03290
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{g}	1.33 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	1405.7875
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^{\pm} \rightarrow q\tilde{q}W^{\pm}\tilde{\chi}_1^0$	0-1 e, μ	2-6 jets	Yes	20	\tilde{g}	1.26 TeV	$m(\tilde{\chi}_1^0)<300 \text{ GeV}, m(\tilde{\chi}_1^{\pm})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$	1507.05525
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20	\tilde{g}	1.32 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	1501.03555
	GMSB ($\tilde{\ell}$ NLSP)	1-2 $\tau + 0-1 \ell$	0-2 jets	Yes	20.3	\tilde{g}	1.6 TeV	$\tan\beta > 20$	1407.0603
	GGM (bino NLSP)	2 γ	-	Yes	20.3	\tilde{g}	1.29 TeV	$c\tau(\text{NLSP})<0.1 \text{ mm}$	1507.05493
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	20.3	\tilde{g}	1.3 TeV	$m(\tilde{\chi}_1^0)<900 \text{ GeV}, c\tau(\text{NLSP})<0.1 \text{ mm}, \mu<0$	1507.05493
GGM (higgsino-bino NLSP)	γ	2 jets	Yes	20.3	\tilde{g}	1.25 TeV	$m(\tilde{\chi}_1^0)<850 \text{ GeV}, c\tau(\text{NLSP})<0.1 \text{ mm}, \mu>0$	1507.05493	
GGM (higgsino NLSP)	2 e, μ (Z)	2 jets	Yes	20.3	\tilde{g}	850 GeV	$m(\text{NLSP})>430 \text{ GeV}$	1503.03290	
Gravitino LSP	0	mono-jet	Yes	20.3	$F^{1/2}$ scale	865 GeV	$m(\tilde{G})>1.8 \times 10^{-4} \text{ eV}, m(\tilde{g})=m(\tilde{q})=1.5 \text{ TeV}$	1502.01518	
3 rd gen. \tilde{g} med.	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	\tilde{g}	1.25 TeV	$m(\tilde{\chi}_1^0)<400 \text{ GeV}$	1407.0600
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	\tilde{g}	1.1 TeV	$m(\tilde{\chi}_1^0)<350 \text{ GeV}$	1308.1841
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^{\pm}$	0-1 e, μ	3 b	Yes	20.1	\tilde{g}	1.34 TeV	$m(\tilde{\chi}_1^0)<400 \text{ GeV}$	1407.0600
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow b\bar{t}\tilde{\chi}_1^{\pm}$	0-1 e, μ	3 b	Yes	20.1	\tilde{g}	1.3 TeV	$m(\tilde{\chi}_1^0)<300 \text{ GeV}$	1407.0600
3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	\tilde{b}_1	100-620 GeV	$m(\tilde{\chi}_1^0)<90 \text{ GeV}$	1308.2631
	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^{\pm}$	2 e, μ (SS)	0-3 b	Yes	20.3	\tilde{b}_1	275-440 GeV	$m(\tilde{\chi}_1^0)=2 m(\tilde{\chi}_1^{\pm})$	1404.2500
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{\chi}_1^{\pm}$	1-2 e, μ	1-2 b	Yes	4.7/20.3	\tilde{t}_1	110-167 GeV	$m(\tilde{\chi}_1^0)=2m(\tilde{\chi}_1^{\pm}), m(\tilde{\chi}_1^0)=55 \text{ GeV}$	1209.2102, 1407.0583
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^0$ or $t\tilde{\chi}_1^0$	0-2 e, μ	0-2 jets/1-2 b	Yes	20.3	\tilde{t}_1	90-191 GeV	$m(\tilde{\chi}_1^0)=1 \text{ GeV}$	1506.08616
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$	0	mono-jet/c-tag	Yes	20.3	\tilde{t}_1	90-240 GeV	$m(\tilde{\chi}_1^0)<85 \text{ GeV}$	1407.0608
	$\tilde{t}_1\tilde{t}_1$ (natural GMSB)	2 e, μ (Z)	1 b	Yes	20.3	\tilde{t}_1	150-580 GeV	$m(\tilde{\chi}_1^0)>150 \text{ GeV}$	1403.5222
	$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$	3 e, μ (Z)	1 b	Yes	20.3	\tilde{t}_2	290-600 GeV	$m(\tilde{\chi}_1^0)<200 \text{ GeV}$	1403.5222
	EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$	90-325 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$
$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$		2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}$	140-465 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^{\pm})+m(\tilde{\chi}_1^0))$	1403.5294
$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\tau}\nu(\tau\bar{\nu})$		2 τ	-	Yes	20.3	$\tilde{\chi}_1^{\pm}$	100-350 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\tau}, \bar{\nu})=0.5(m(\tilde{\chi}_1^{\pm})+m(\tilde{\chi}_1^0))$	1407.0350
$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_1\nu\tilde{\ell}_1\ell(\bar{\nu}\nu), (\bar{\nu}\tilde{\ell}_1\ell(\bar{\nu}\nu))$		3 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	700 GeV	$m(\tilde{\chi}_1^0)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{\chi}_1^0))$	1402.7029
$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$		2-3 e, μ	0-2 jets	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	420 GeV	$m(\tilde{\chi}_1^0)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{ sleptons decoupled}$	1403.5294, 1402.7029
$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 h\tilde{\chi}_1^0, h \rightarrow b\bar{b}/WW/\tau\tau/\gamma\gamma$		e, μ, γ	0-2 b	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	250 GeV	$m(\tilde{\chi}_1^0)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{ sleptons decoupled}$	1501.07110
$\tilde{\chi}_2^0\tilde{\chi}_3^0, \tilde{\chi}_2^0 \rightarrow \tilde{\ell}_R\ell$		4 e, μ	0	Yes	20.3	$\tilde{\chi}_{2,3}^0$	620 GeV	$m(\tilde{\chi}_2^0)=m(\tilde{\chi}_3^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_2^0)+m(\tilde{\chi}_1^0))$	1405.5086
GGM (wino NLSP) weak prod.		1 $e, \mu + \gamma$	-	Yes	20.3	\tilde{W}	124-361 GeV	$c\tau<1 \text{ mm}$	1507.05493
Long-lived particles		Direct $\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$ prod., long-lived $\tilde{\chi}_1^{\pm}$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^{\pm}$	270 GeV	$m(\tilde{\chi}_1^0)-m(\tilde{\chi}_1^{\pm})\sim 160 \text{ MeV}, \tau(\tilde{\chi}_1^{\pm})=0.2 \text{ ns}$
	Direct $\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$ prod., long-lived $\tilde{\chi}_1^{\pm}$	dE/dx trk	-	Yes	18.4	$\tilde{\chi}_1^{\pm}$	482 GeV	$m(\tilde{\chi}_1^0)-m(\tilde{\chi}_1^{\pm})\sim 160 \text{ MeV}, \tau(\tilde{\chi}_1^{\pm})<15 \text{ ns}$	1506.05332
	Stable, stopped \tilde{g} R-hadron	0	1-5 jets	Yes	27.9	\tilde{g}	832 GeV	$m(\tilde{\chi}_1^0)=100 \text{ GeV}, 10 \mu\text{s}<\tau(\tilde{g})<1000 \text{ s}$	1310.6584
	Stable \tilde{g} R-hadron	trk	-	-	19.1	\tilde{g}	1.27 TeV		1411.6795
	GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{\ell}, \bar{\mu})+\tau(e, \mu)$	1-2 μ	-	-	19.1	$\tilde{\chi}_1^0$	537 GeV	$10<\tan\beta<50$	1411.6795
	GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	-	Yes	20.3	$\tilde{\chi}_1^0$	435 GeV	$2<\tau(\tilde{\chi}_1^0)<3 \text{ ns}, \text{SPS8 model}$	1409.5542
	$\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow e\bar{e}\nu/\mu\bar{\mu}\nu/\mu\bar{\mu}\nu$	displ. $e\bar{e}/\mu\bar{\mu}/\mu\bar{\mu}\nu$	-	-	20.3	$\tilde{\chi}_1^0$	1.0 TeV	$7<c\tau(\tilde{\chi}_1^0)<740 \text{ mm}, m(\tilde{g})=1.3 \text{ TeV}$	1504.05162
	GGM $\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow Z\tilde{G}$	displ. vtx + jets	-	-	20.3	$\tilde{\chi}_1^0$	1.0 TeV	$6<c\tau(\tilde{\chi}_1^0)<480 \text{ mm}, m(\tilde{g})=1.1 \text{ TeV}$	1504.05162
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e\mu/\tau\mu/\mu\tau$	$e\mu, e\tau, \mu\tau$	-	-	20.3	$\tilde{\nu}_\tau$	1.7 TeV	$\lambda'_{341}=0.11, \lambda_{132/133/233}=0.07$	1503.04430
	Bilinear RPV CMSSM	2 e, μ (SS)	0-3 b	Yes	20.3	\tilde{q}, \tilde{g}	1.35 TeV	$m(\tilde{q})=m(\tilde{g}), c\tau_{LS}\mu<1 \text{ mm}$	1404.2500
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow e\bar{e}\nu_{\mu}, e\mu\bar{\nu}_e$	4 e, μ	-	Yes	20.3	$\tilde{\chi}_1^{\pm}$	750 GeV	$m(\tilde{\chi}_1^0)>0.2 \times m(\tilde{\chi}_1^{\pm}), \lambda_{121}\neq 0$	1405.5086
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tau\tau\nu_e, e\tau\bar{\nu}_e$	3 $e, \mu + \tau$	-	Yes	20.3	$\tilde{\chi}_1^{\pm}$	450 GeV	$m(\tilde{\chi}_1^0)>0.2 \times m(\tilde{\chi}_1^{\pm}), \lambda_{133}\neq 0$	1405.5086
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}q$	0	6-7 jets	-	20.3	\tilde{g}	917 GeV	$\text{BR}(\mu)=\text{BR}(b)=\text{BR}(c)=0\%$	1502.05686
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow q\tilde{q}q$	0	6-7 jets	-	20.3	\tilde{g}	870 GeV	$m(\tilde{\chi}_1^0)=600 \text{ GeV}$	1502.05686
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t_1\bar{t}, \bar{t}_1 \rightarrow b\bar{s}$	2 e, μ (SS)	0-3 b	Yes	20.3	\tilde{g}	850 GeV		1404.250
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\bar{s}$	0	2 jets + 2 b	-	20.3	\tilde{t}_1	100-308 GeV		ATLAS-CONF-2015-026
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\bar{\ell}$	2 e, μ	2 b	-	20.3	\tilde{t}_1	0.4-1.0 TeV	$\text{BR}(\tilde{t}_1 \rightarrow b\bar{\ell}/\mu)>20\%$	ATLAS-CONF-2015-015
Other	Scalar charm, $\tilde{c} \rightarrow c\tilde{\chi}_1^0$	0	2 c	Yes	20.3	\tilde{c}	490 GeV	$m(\tilde{\chi}_1^0)<200 \text{ GeV}$	1501.01325

10^{-1}

1

Mass scale [TeV]

10^{-1}

1

Mass scale [TeV]




*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.

And even ...

... a little intrigue

CMS-PAS-EXO-15-004


ATLAS-CONF-2015-081



To the Future!

LHC → HL-LHC


<http://hilumilhc.web.cern.ch/about/hl-lhc-project>



today: Higgs Maxwell meeting 2016

The Challenge of Pileup

- Pileup = number of proton-proton collision per bunch crossing
- Instantaneous luminosity of $5\text{ (7)} \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ corresponds to an average pileup of $\langle\mu\rangle$ of 140 (200).




Simulated pileup in ATLAS tracker

Run 1

Pile up of 23

HL-HLC
Pile up of 230



ATLAS and CMS Upgrades

CERN-LHC


CERN-LHC

- ATLAS and CMS will be upgraded to achieve the same or better performance as in Run 1.
 - ▶ Pileup mitigation is a critical element of detector designs.
- Recently released detector *Scoping Documents* investigate the impact of different detector cost scenarios on physics performance.
- e.g. for 2022: New tracking detectors, new trigger systems, new timing detectors.

ATLAS



CMS




HL-LHC Analysis Techniques

High Pileup

- Much effort is focussed on understanding how to mitigate pileup in physics analyses
- e.g. New method proposed in the literature *Pileup Per Particle Identification* [arXiv:1407.6013](https://arxiv.org/abs/1407.6013)


ATLAS:

Resolution as a function of ΣE_T in $t\bar{t}$ events:
use extended tracking to reject pile-up jets



CMS: Rate of pileup jets/true jets for


- Particle Flow algorithm (PF)
- PF + rejecting charged hadrons from pileup
- Using Puppi algorithm




HL-LHC Analysis Techniques

Jet Substructure

- High mass final states and high collision energy lead to highly boosted and close objects e.g. $W \rightarrow jj$, $Z \rightarrow jj$, $t \rightarrow Wb \rightarrow jjb$
- Jet substructure techniques will be key to reconstruct some of these signals; crucial for new high-mass objects.



HL-LHC Physics Prospects






- Results are projections from refining current analyses or designing new ones.
- In some cases, several different systematic uncertainty scenarios are presented.
- Many results are presented in the context of specific models.

Higgs Boson Physics

- HL-LHC will be a Higgs boson factory \Rightarrow over 100 million Higgs bosons in 3000 fb^{-1}

➔ Over 1 million for each of the main production mechanisms, spread over many decay modes

- 400k $H \rightarrow \gamma\gamma$
- 20k $H \rightarrow ZZ \rightarrow llll$
- 40k $H \rightarrow \mu\mu$
- 50 leptonic $H \rightarrow J/\psi \gamma$ (very rare mode)




Higgs Boson Peaks with 3000 fb⁻¹

ATL-PHYS-PUB-2013-014


ATL-PHYS-PUB-2014-012

CERN-LHCC-2015-010


VBF, $H \rightarrow WW$




ZH, $H \rightarrow b\bar{b}$



**$t\bar{t}H, H \rightarrow \gamma\gamma$
1 lepton**



$H \rightarrow ZZ \rightarrow \ell\ell\ell\ell$



VBF $H \rightarrow WW$

- Used to motivate the ATLAS upgrade detector design in the Scoping Document.
- Two forward jets in the detector

Mass of two forward jets:










Table 35. The $\Delta\mu/\mu$ and significance for VBF $H \rightarrow WW^{(*)}$ are shown for the three scoping scenarios. Results with and without the theoretical uncertainties on the VBF or ggF Higgs boson production are included.

Scoping Scenario	without theo. unc.		with theo. unc.	
	$\Delta\mu/\mu$	Z_0 -value (σ)	$\Delta\mu/\mu$	Z_0 -value (σ)
Reference	0.14	8.0	0.20	5.7
Middle	0.20	5.4	0.25	4.4
Low	0.30	3.5	0.39	2.7

Still Golden: $H \rightarrow ZZ \rightarrow \ell\ell\ell\ell$

ATL-PHYS-PUB-2013-014


signal events	ggH	VBF	$t\bar{t}H$	WH	ZH
3000 fb^{-1}	3800	97	35	67	6



Large statistics will be used for $d\sigma/dp_T(H)$, $d\sigma/dN_{\text{jets}}$

Higgs CP Studies

- $H \rightarrow ZZ \rightarrow 4\ell$ used to reconstruct the full angular decay structure.
- Very sensitive to non-SM ($\mathbf{CP} = 0^+$) contributions.



$$A(H \rightarrow ZZ) = v^{-1} \left(\underbrace{a_1 m_Z^2 \epsilon_1^* \epsilon_2^*}_{\text{SM tree processes}} + \underbrace{a_2 f_{\mu\nu}^{*(1)} f^{*(2),\mu\nu}}_{\text{loop CP-even contributions}} + \underbrace{a_3 f_{\mu\nu}^{*(1)} \tilde{f}^{*(2),\mu\nu}}_{\text{CP-odd contributions (BSM)}} \right)$$


- Fit fraction of event (f_{ai}) and phases (ϕ_i) to observed decay:

$$\phi_{a_i} = \arg \left(\frac{a_i}{a_1} \right) \quad f_{a_i} = \frac{|a_i|^2 \sigma_i}{|a_1|^2 \sigma_1 + |a_i|^2 \sigma_i}$$

Higgs CP Studies


arXiv:1307.7135

ATL-PHYS-PUB-2013-013



Loop-induced CP-even contribution

$2 \times \Delta(\text{NLL})$




CP-odd contribution

- Extra contributions constrained to $|f| \sim 10\%$ with 3000 fb $^{-1}$.



Rare Processes

ATL-PHYS-PUB-2013-014

- $H \rightarrow \mu\mu$ - measures coupling to second fermion generation
 - ▶ ATLAS and CMS expect $>7\sigma$ significance with 3000 fb^{-1}
 - ▶ coupling measured to 5-10%



- $H \rightarrow Z\gamma$
 - ▶ Tests loop structure of decay, compare with $H \rightarrow ZZ$ $H \rightarrow \gamma\gamma$
 - ▶ $\sim 4\sigma$ significance possible with 3000 fb^{-1} despite the challenging background



Higgs Boson Signal Strength Sensitivity


ATL-PHYS-PUB-2014-016

arXiv:1307.7135

- All production modes can be observed for ZZ and $\gamma\gamma$ final states
- Combine production modes for best information on branching ratios

ATLAS Simulation Preliminary


$\sqrt{s} = 14$ TeV: $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$; $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$



dashed bands
includes
current
theory
uncertainties

Scenario 2:
TH unc. scaled by 1/2
EXP unc. scaled by $\sqrt{\mathcal{L}}$

CMS Projection



3000 fb^{-1}


Summary of precision for 3000 fb^{-1} :

- ~ 4 - 5% for main channels
- ~ 10 - 20% on rare modes

Higgs Boson Width

ATL-PHYS-PUB-2015-024

- $H \rightarrow ZZ \rightarrow 4\ell$: Use the interference between off-shell and on-shell production to measure $\Gamma(H)$




- For 3000 fb^{-1} ; using uncertainty between background signal of $\sigma(R_{H^*}^B) = 10\%$; combining on-shell and off-shell measurement; assuming off-shell measurement dominates, for $\Gamma = \Gamma_{\text{SM}}$ gives:

$$\Gamma_H = 4.2^{+1.5}_{-2.1} \text{ MeV (stat+sys)}$$

Interpretation as Coupling Scale Factors

- Experiments measure cross section times branching ratio
- Interpretation with coupling scale factors, κ , is model dependent



• e.g.


$$(\sigma \cdot \text{BR})(gg \rightarrow H \rightarrow \gamma\gamma) = \sigma_{\text{SM}}(gg \rightarrow H) \cdot \text{BR}_{\text{SM}}(H \rightarrow \gamma\gamma) \cdot \frac{\kappa_g^2 \cdot \kappa_\gamma^2}{\kappa_H^2}$$

- Assume Γ_H is sum of sum of visible widths - no additional invisible modes


Higgs Boson Couplings Fit

arXiv:1307.7135
ATL-PHYS-PUB-2014-016

CMS Projection




CMS Projection



ATLAS Simulation Preliminary

$\sqrt{s} = 14$ TeV: $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$; $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$



Mass scaled couplings



CERN-LHCC-2015-010
ATL-PHYS-PUB-2014-016

Vector boson coupling

$$y_{V,i} = \sqrt{\kappa_{V,i} \frac{g_{V,i}}{2v}} = \sqrt{\kappa_{V,i}} \frac{m_{V,i}}{v}$$

Fermion couplings

$$y_{f,i} = \kappa_{f,i} \frac{g_{f,i}}{\sqrt{2}} = \kappa_{f,i} \frac{m_{f,i}}{v}$$





HH




Higgs Boson Pair Production

- Higgs boson pair production includes destructive interference between two types of processes:



[arXiv:1401.7340v2](https://arxiv.org/abs/1401.7340)

NNLO $\sigma^{\text{SM}} = 40.8 \text{ fb}$




Number of events in 3000 fb ⁻¹	
<i>bbWW</i>	30000
<i>bbττ</i>	9000
<i>WWWW</i>	6000
<i>γγbb</i>	320
<i>γγγγ</i>	1



e.g. $HH \rightarrow bb\gamma\gamma$

CERN-LHCC-2015-010
ATL-PHYS-PUB-2014-019

- bb mass peak is broad
- $\gamma\gamma$ shows narrow resonance




- **CMS:** 2d fit of $m(bb)$ and $m(\gamma\gamma)$ distributions (control background from data)
- **ATLAS** cut based analysis



$H \rightarrow \gamma\gamma, H \rightarrow b\bar{b}$ candidate event at $\sqrt{s}=8$ TeV

[arXiv:1406.5053](https://arxiv.org/abs/1406.5053)







$HH \rightarrow bb\tau\tau$

CERN-LHCC-2015-010

ATL-PHYS-PUB-2015-046

- Major background from $t\bar{t}$, with $t \rightarrow \tau\nu b$
- Results for 3000 fb^{-1} and SM ($\lambda=1$):
 - ATLAS: All $\tau\tau$ final states considered
 - ▶ **13 $HH \rightarrow bb\tau\tau$ events** after full event selection cf 803 background events
 - CMS: $\tau(\text{had}) \tau(\text{had})$ & $\tau(\mu) \tau(\text{had})$; kinematic variables to distinguish signal from background
 - ▶ Just 0.9σ sensitivity




Vector Boson Scattering

Vector Boson Scattering



[arXiv:1405.6241](#)

[arXiv:1410.6315](#)

- Explore electroweak symmetry breaking through VBS: e.g. Look for W^+W^+ , W^-W^- and WZ final states




- In Run 1, ATLAS (CMS) observed 4.5σ (2.0σ) evidence for $W^\pm W^\pm jj$ production
- CMS: Limits placed on dimension-8 operators, f_X/Λ^4 (à la Eboli, Gonzalez-Garcia, Mizukoshi [arXiv:hep-ph/0606118](#))





Run 1 Evidence for Weak Boson Scattering

[arXiv:1405.6241](https://arxiv.org/abs/1405.6241)



Weak Boson Scattering Prospects


- WW : Very clear signature expected with sensitivity to 125 GeV Higgs boson propagator
- WZ : More challenging due to large QCD contribution.



Beyond the Standard Model

Additional Heavy Higgs bosons



- Additional Higgs doublets predicted in many models, including Supersymmetry.
- e.g. A two-Higgs doublet (2HDM) model includes four new Higgs boson:



$\cos(\beta-\alpha) \rightarrow 0$
if h^0 is SM-like


- α is a mixing angle between the Higgs doublets
- $\tan\beta$ is the ratio between the vev the Higgs doublets

$A \rightarrow Zh \rightarrow \ell\ell b\bar{b}$ reconstruction (2HDM)




Higgs Portal to Dark Matter

- In the SM Higgs boson couples to all massive particles
 - ▶ very likely Higgs boson will also couple to any DM WIMPs, χ
 - ▶ look for a branching ratio for Higgs boson to invisible particles
 - ▶ Coupling of χ to H take as free parameter; **BR(inv)** sets a limit on the interactions of χ
- In 3000 fb^{-1}
 - ▶ **ATLAS: $\text{BR(inv)} < 0.13$**
(0.09 w/out theory uncertainties)
 - ▶ **CMS: $\text{BR(inv)} < 0.11$**
(0.07 in alt. theory uncertainty)



LHC complements direct DM search experiments in the lower mass range

SUSY production at the LHC



The lightest neutralino (LSP) is candidate to explain dark matter.


Stop and Sbottom Searches

CMS-PAS-FTR-13-014


ATL-PHYS-PUB-2013-011

ATL-PHYS-PUB-2014-010

Stop pair production; $\tilde{t} \rightarrow t\tilde{\chi}_1^0$



Sbottom pair production; $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$




5σ discovery, simplified model	300 fb ⁻¹	3000 fb ⁻¹
stop mass from direct production [ATLAS]	Up to 1.0 TeV	Up to 1.2 TeV
gluino mass with decay to stop [CMS]	Up to 1.9 TeV	Up to 2.2 TeV
sbottom mass from direct production [ATLAS]	Up to 1.1 TeV	Up to 1.3 TeV

Strong and Weak SUSY Production Limits

CMS-PAS-FTR-13-014
ATL-PHYS-PUB-2014-010

Strong SUSY: Gluino pair production




Large uncertainties on σ from knowledge of PDFs

Weak SUSY: Chargino and neutralino decaying via WZ

$$\chi_1^\pm \rightarrow W^\pm \chi_1^0,$$

$$\chi_2^0 \rightarrow Z \chi_1^0$$



Simplified SUSY model

Chargino mass 5 σ discovery, simplified model		300 fb ⁻¹	3000 fb ⁻¹
WZ (3l analysis)	[ATLAS]	Up to 560 GeV	Up to 820 GeV
WZ (3l analysis)	[CMS]	Up to 600 GeV	Up to 900 GeV
WH (3l analysis)	[ATLAS]	(<5 σ reach)	Up to 650 GeV
WH (bb analysis)	[ATLAS] (new in 2015)	(<5 σ reach)	Up to 800 GeV
WH (bb analysis)	[CMS]	350-460 GeV	Up to 950 GeV

Summary of SUSY Simplified Models Reach

ATLAS projection	gluino mass	squark mass	stop mass	sbottom mass	χ_1^+ mass WZ mode	χ_1^+ mass WH mode
300 fb ⁻¹	2.0 TeV	2.6 TeV	1.0 TeV	1.1 TeV	560 GeV	None
3000 fb ⁻¹	2.4 TeV	3.1 TeV	1.2 TeV	1.3 TeV	820 GeV	650 GeV

Resonance Searches

arXiv:1307.7135


ATL-PHYS-PUB-2014-007

ATL-PHYS-PUB-2013-003


- New physics could appear anywhere!

- Look for resonances in di-leptons, $\gamma\gamma$, $t\bar{t}$, di-bosons (WW , WZ , ZZ) and extra missing transverse momentum.


4 TeV Kaluza-Klein gluon, $g_{KK} \rightarrow t\bar{t}$




5 TeV $Z' \rightarrow \mu^+\mu^-$



$Z' \rightarrow e^+e^-$



Mass Reach for Exotic Signatures



ATLAS @14 TeV	$Z' \rightarrow ee$ SSM 95% CL limit	$g_{KK} \rightarrow tt$ RS 95% CL limit	Dark matter M^* 5 σ discovery
300 fb ⁻¹	6.5 TeV	4.3 TeV	2.2 TeV
3000 fb ⁻¹	7.8 TeV	6.7 TeV	2.6 TeV

Top Quark Physics



Top Quark Physics

ATL-PHYS-PUB-2013-012

CMS-PAS-FTR-13-016

CMS-PAS-FTR-13-017

- *Top quark mass parameter* can be measured to $\Lambda_{\text{QCD}} \sim 200 \text{ MeV}$ in 3000 fb^{-1} .
 - ▶ Endpoint method, which probes the pole mass, can measure m_t to 500 MeV
- In SM $\text{BR}(t \rightarrow Wb) \approx 100\%$ Many models predict enhancements, interesting range starts at $\sim 10^{-4} \Rightarrow$ Observing decays to other modes clear sign of new physics
 - ▶ HL-LHC will probe $\text{BR}(t \rightarrow qZ)$, $\text{BR}(t \rightarrow q\gamma)$ at $\sim 3 \times 10^{-5}$ at least and $\text{BR}(t \rightarrow cH)$ at $\sim 10^{-4}$



Outlook

- We've come a long way, baby, but there's still far to go...
- With 3000 fb^{-1} the LHC will offer a comprehensive physics programme:

Precision Higgs physics:
measure production rates
to a few % (model dep.)

SUSY: Assuming light LSP ($<1 \text{ TeV}$)
discover squarks up to 1.1 TeV
discover gluinos up to 2 TeV

Sensitivity to generic
resonances and missing
energy up to $O(7 \text{ TeV})$

Observation of
 $H \rightarrow Z\gamma$ and $H \rightarrow \mu^+\mu^-$

Measure m_{top} to 200 MeV
Sensitivity to rare top
quark decays of $<10^{-4}$

HH observation ...
might reach 5σ

Discovery of additional Higgs
bosons up to $O(1 \text{ TeV})$

Theory uncertainty dominant for many analyses

- Some analyses do remain beyond the reach of HL-LHC:

$H \rightarrow c\bar{c}$


triple-Higgs boson



CMS Experiment at the LHC, CERN

Data recorded: 2015-Sep-28 06:09:43.129280 GMT

Run / Event / LS: 257645 / 1610868539 / 1073



CMS's highest mass event
12 jet with $p_T > 50$ GeV each!
Total mass of system 6.4 TeV

Backup

ATLAS Upgrades

- Long Shutdown 1


- New beam pipe at $r=25\text{mm}$
 - New insertable b -layer at $31 < r/\text{mm} < 40$
 - Refurbished pixel readout
 - More complete muon coverage: extended endcap installation complete
- Fast Tracking for L2-trigger will come online during run 2

- Long Shutdown 2

- New muon small wheel forward spectrometer
- Topological L1-trigger processors
- New forward detectors

- For HL-LHC

- Completely new trigger architecture with new hardware at L0/L1
- Completely new tracking detector
- 53 ● Calorimeter electronics upgrades



ATLAS & CMS Higgs boson coupling fits

L(fb ⁻¹)	Exp.	κ_γ	κ_W	κ_Z	κ_g	κ_b	κ_t	κ_τ	$\kappa_{Z\gamma}$	$\kappa_{\mu\mu}$
300	ATLAS	[9, 9]	[9, 9]	[8, 8]	[11, 14]	[22, 23]	[20, 22]	[13, 14]	[24, 24]	[21, 21]
	CMS	[5, 7]	[4, 6]	[4, 6]	[6, 8]	[10, 13]	[14, 15]	[6, 8]	[41, 41]	[23, 23]
3000	ATLAS	[4, 5]	[4, 5]	[4, 4]	[5, 9]	[10, 12]	[8, 11]	[9, 10]	[14, 14]	[7, 8]
	CMS	[2, 5]	[2, 5]	[2, 4]	[3, 5]	[4, 7]	[7, 10]	[2, 5]	[10, 12]	[8, 8]

- Long Shutdown 1:


- Complete Muon coverage
- New HCAL photo-detectors

- Long Shutdown 2:


- New Pixel detector (2017)
- New HCAL electronics
- L1-Trigger upgrade

- For HL-LHC:

- Tracker replacement, L1 Track-Trigger
- New forward calorimetry, muons and tracking
- High precision timing for pileup mitigation





Greater trigger efficiency



Run 1 Top Quark Properties


[arXiv:1403.4427](https://arxiv.org/abs/1403.4427)

[CMS-PAS-TOP-14-015](#)



Run 1 observed limits on stop and LSP

[arXiv:1506.08616](https://arxiv.org/abs/1506.08616)





Simplified models
with $\tilde{t} \rightarrow \text{LSP} + X$

HL-LHC: Additional Heavy Higgs bosons


Prospects for $H' \rightarrow ZZ \rightarrow 4\ell$ production

ATL-PHYS-PUB-2013-016

CMS-PAS-FTR-13-024




Prospects for $\phi \rightarrow \mu\mu$ production



Run 1 SUSY limits

[arXiv:1507.05525](https://arxiv.org/abs/1507.05525)



HL-LHC Cross Sections

proton - (anti)proton cross sections

