

## Formation and Stability of Dense Methane-Hydrogen Compounds

Umbertoluca Ranieri<sup>1,2</sup>, Lewis J. Conway<sup>3</sup>, Mary-Ellen Donnelly<sup>1</sup>, Huixin Hu<sup>1</sup>, Mengnan Wang<sup>1</sup>, Philip Dalladay-Simpson<sup>1</sup>, Miriam Peña-Alvarez<sup>3</sup>, Eugene Gregoryanz<sup>1,3,4</sup>, Andreas Hermann<sup>3</sup>, and Ross T. Howie<sup>1,3,\*</sup>

<sup>1</sup>Center for High Pressure Science and Technology Advanced Research, 1690 Cailun Road, Shanghai, 201203, China

<sup>2</sup>Dipartimento di Fisica, Università di Roma La Sapienza, Piazzale Aldo Moro 5, 00185 Rome, Italy

<sup>3</sup>Centre for Science at Extreme Conditions and The School of Physics and Astronomy, The University of Edinburgh, Peter Guthrie Tait Road, Edinburgh, United Kingdom

<sup>4</sup>Key Laboratory of Materials Physics, Institute of Solid State Physics, CAS, Hefei, China

 (Received 25 November 2021; revised 2 February 2022; accepted 20 April 2022; published 27 May 2022)

Through a series of x-ray diffraction, optical spectroscopy diamond anvil cell experiments, combined with density functional theory calculations, we explore the dense CH<sub>4</sub>-H<sub>2</sub> system. We find that pressures as low as 4.8 GPa can stabilize CH<sub>4</sub>(H<sub>2</sub>)<sub>2</sub> and (CH<sub>4</sub>)<sub>2</sub>H<sub>2</sub>, with the latter exhibiting extreme hardening of the intramolecular vibrational mode of H<sub>2</sub> units within the structure. On further compression, a unique structural composition, (CH<sub>4</sub>)<sub>3</sub>(H<sub>2</sub>)<sub>25</sub>, emerges. This novel structure holds a vast amount of molecular hydrogen and represents the first compound to surpass 50 wt % H<sub>2</sub>. These compounds, stabilized by nuclear quantum effects, persist over a broad pressure regime, exceeding 160 GPa.

DOI: [10.1103/PhysRevLett.128.215702](https://doi.org/10.1103/PhysRevLett.128.215702)

Hydrogen and methane are, besides water, the most prevalent small molecules in the outer Solar System. Their interplay under extreme conditions is of key interest to understanding the evolution and interior dynamics of Neptune and Uranus, as well as of Earth and exoplanets [1–4]. The simplest molecule, H<sub>2</sub>, has been shown to exhibit rich physical phenomena, including quantum rotational phases and pressure-induced steps toward an atomic metallic solid, while methane (CH<sub>4</sub>), the simplest hydrocarbon, has been proposed to polymerize during compression to form long-chain hydrocarbons and at further extremes decompose into diamond plus H<sub>2</sub> [5–19].

Hydrogen reacts with a number of materials at high pressures and temperatures but also has a propensity to form van der Waals compounds that can be stable far beyond 100 GPa [20–32]. Methane and hydrogen were first reported to crystallize into inclusion (host-guest) compounds over 25 years ago at pressures between 5 and 7 GPa. A range of compositions were claimed: (CH<sub>4</sub>)<sub>2</sub>H<sub>2</sub>, CH<sub>4</sub>H<sub>2</sub>, CH<sub>4</sub>(H<sub>2</sub>)<sub>2</sub>, and CH<sub>4</sub>(H<sub>2</sub>)<sub>4</sub> [21]. However, the experimental evidence for these compounds was limited, with only the shift of the hydrogen intramolecular vibrational mode  $\nu_1$ -H<sub>2</sub> (vibron) relative to that of pure hydrogen being reported together with the lattice parameters of potential structures. Assuming that these structural compositions were correct, a later experimental study explored the recoverability of CH<sub>4</sub>(H<sub>2</sub>)<sub>4</sub> at low temperature to investigate its potential as a hydrogen storage media [33,34].

In contrast to experimental work, the CH<sub>4</sub>-H<sub>2</sub> system has recently had a surge of theoretical investigations, predicting a variety of extremely hydrogen-rich compositions to emerge such as triclinic (CH<sub>4</sub>)<sub>2</sub>(H<sub>2</sub>)<sub>3</sub>, trigonal (CH<sub>4</sub>)<sub>2</sub>(H<sub>2</sub>)<sub>7</sub>, and

hexagonal CH<sub>4</sub>(H<sub>2</sub>)<sub>9</sub>, as well as several CH<sub>4</sub>H<sub>2</sub> structures [18,35–37]. Furthermore, it was suggested that CH<sub>4</sub>-H<sub>2</sub> structures could be the most stable form of carbon and hydrogen at pressures up to 200 GPa [18].

Surprisingly, given these predictions and the advent of technological advances over the past 25 years, no further experimental exploration has been conducted despite undetermined CH<sub>4</sub>-H<sub>2</sub> compound signatures emerging as a by-product in many prolific studies of hydrocarbons at planetary conditions [3,19,38–41]. Moreover, the doping of carbon (or methane) could even enhance the properties of materials at extreme pressures, with the unprecedented claim of room temperature superconductivity in the carbonaceous sulfur hydride system [42–44]. The lack of knowledge regarding CH<sub>4</sub>-H<sub>2</sub> compound formation inhibits our understanding of these more complex ternary systems and how doping could induce novel properties. As such, it is of fundamental interest to know which methane-hydrogen compounds are actually formed and test their pressure stability limits.

In this Letter, we have conducted a series of high-pressure synchrotron powder x-ray diffraction and Raman spectroscopy experiments in diamond anvil cells combined with density functional theory (DFT) calculations and structural searching to explore the formation and properties of CH<sub>4</sub>-H<sub>2</sub> compounds from mixtures of methane and hydrogen. In H<sub>2</sub>-rich mixtures, we observe the formation of hexagonal CH<sub>4</sub>(H<sub>2</sub>)<sub>2</sub> above 5 GPa, before partially transforming into monoclinic (CH<sub>4</sub>)<sub>3</sub>(H<sub>2</sub>)<sub>25</sub>. (CH<sub>4</sub>)<sub>3</sub>(H<sub>2</sub>)<sub>25</sub> represents a unique composition and contains an unprecedented 51.1 wt % H<sub>2</sub>, which is the highest hydrogen content of any currently known




FIG. 1. Pressure-composition phase diagram of the  $\text{CH}_4\text{-H}_2$  binary system. Filled square symbols indicate the formation pressures of each compound:  $(\text{CH}_4)_2\text{H}_2$  (orange),  $\text{CH}_4(\text{H}_2)_2$  (purple), and  $(\text{CH}_4)_3(\text{H}_2)_{25}$  (green). The green open squares represent the pressures at which  $(\text{CH}_4)_3(\text{H}_2)_{25}$  is first observed. The black squares, black triangle, and blue circle represent the formation pressure of  $\text{CH}_4\text{-I}$ ,  $\text{CH}_4\text{-A}$ , and  $\text{H}_2\text{-I}$ , respectively. The gray line represents the liquidus curve and is adapted from Ref. [21]. Below 7 GPa, the error in pressure is  $\pm 0.2$  GPa and smaller than the size of the symbol. Initial gas mixtures have a tolerance of 1%.

stoichiometric compound. In  $\text{CH}_4$ -rich mixtures, tetragonal  $(\text{CH}_4)_2\text{H}_2$  forms, which undergoes extreme hardening of the  $\text{H}_2$  intramolecular vibrational mode with pressure. All three compounds exhibit remarkable stability, exceeding pressures of 160 GPa.

At pressures below 1.5 GPa, all mixture concentrations (30, 50, 70, 80, and 90 mol %  $\text{H}_2$ ) are homogeneous well-mixed fluids and display the Raman signatures of both constituent species (see Fig. 1 for the pressure-composition phase diagram, Supplemental Material [45] for experimental methods, and Figs. S1–S9 [45] for Raman spectra of all mixtures). On compression of hydrogen-rich  $\text{CH}_4\text{-H}_2$  mixtures (70–90 mol %  $\text{H}_2$ ) above 5 GPa, the fluid mixture crystallizes into a solid which we identify as  $\text{CH}_4(\text{H}_2)_2$ . X-ray diffraction measurements reveal that this compound adopts a hexagonal  $\text{MgZn}_2$  Laves phase structure (space group  $P6_3/mmc$ ), where the  $\text{CH}_4$  molecules occupy the Mg sites and the  $\text{H}_2$  molecules occupy the Zn sites (see Fig. 2), the lattice parameters of which are  $a = 4.981$  Å and  $c = 8.125$  Å at 16.6 GPa. This structure and composition was suggested previously but attributed to a compound forming in a narrow pressure regime (between approximately 6 and 7 GPa) and between 35 and 65 mol %  $\text{H}_2$  mixture concentrations [21]. Here, we find that  $\text{CH}_4(\text{H}_2)_2$  is the dominant phase across all  $\text{H}_2$ -rich mixtures and persistent over a large pressure regime. Raman spectroscopy

reveals two distinct  $\text{H}_2$  vibrons,  $\nu_{1H}$  and  $\nu_{2H}$ , the latter of which is approximately 50 times lower in intensity. The two most intense  $\text{CH}_4$  stretching modes,  $\nu_{1M}$  and  $\nu_{2M}$ , are shifted to higher frequency compared to pure  $\text{CH}_4$ , while the wagging mode is shifted to lower frequency (see Figs. 3 and 4).

On further compression, we observe another compound emerge evidenced by visual changes in the sample morphology and the appearance of three distinct  $\text{H}_2$  vibrational modes (the lowest in frequency being the most intense by 2 orders of magnitude) and  $\text{CH}_4$  stretching bands that can be isolated from  $\text{CH}_4(\text{H}_2)_2$  (see Figs. 3 and 4 and Supplemental Material [45]). The most intense  $\text{H}_2$  vibron is lower in frequency than that of  $\text{CH}_4(\text{H}_2)_2$ , suggestive that the compound has a higher  $\text{H}_2$  content. X-ray diffraction measurements indicate the compound is weakly scattering; however, this could be in part due to only partial transformation of the sample. The patterns could be indexed to a hexagonal cell, with  $a = 7.804$  Å and  $c = 11.199$  Å at 17.3 GPa (see Fig. 2). Systematic absences, unit cell volume, and  $c/a$  ratio suggest this compound has a close resemblance with the previously reported  $R\bar{3}m$  structure of  $\text{Xe}(\text{H}_2)_8$  [23,26] and an initial composition estimate of  $\text{CH}_4(\text{H}_2)_8$ . This seemed plausible given  $\text{CH}_4$  and  $\text{Xe}$  have similar van der Waals diameters (3.78 and 4.32 Å, respectively).

We subsequently performed DFT calculations to help identify this phase. We constructed a series of structures of variable composition, starting by populating the hexagonal cell with randomly oriented  $\text{CH}_4$  molecules on the Xe sites of  $\text{Xe}(\text{H}_2)_8$ , before adding  $\text{H}_2$  molecules on hexagonal close packed (hcp) arrangements commensurate with the cell, using up to  $3 \times 3 \times 4$  repeats of the hcp lattice and a random offset against the  $\text{CH}_4$  sublattice.  $\text{H}_2$  molecules too close to  $\text{CH}_4$  were removed before fully optimizing the remaining atoms and all lattice parameters. Several thousand of these structures were generated with stoichiometries ranging from  $\text{CH}_4(\text{H}_2)_7$  to  $\text{CH}_4(\text{H}_2)_{11}$ . Of these,  $\text{CH}_4(\text{H}_2)_{8.33}$  [or  $(\text{CH}_4)_3(\text{H}_2)_{25}$ ] emerged as the most energetically competitive and was found to be dynamically stable at and above 20 GPa (see Figs. S20 and S21 [45]). It is also the only computationally obtained structure which reproduces the positions and intensities of the experimental diffraction peaks across the studied pressure range (see Fig. S14 [45]), which is meaningful given that no symmetry restrictions were applied in the structures' construction and subsequent optimization. While we find both  $\text{CH}_4(\text{H}_2)_2$  and  $(\text{CH}_4)_2\text{H}_2$  stable or close to stability within the  $\text{CH}_4\text{-H}_2$  phase diagram across a wide pressure range, once zero point energies and vibrational entropies are accounted for,  $(\text{CH}_4)_3(\text{H}_2)_{25}$  remains metastable throughout (see Supplemental Material [45]).

Considering methane and hydrogen molecules to be spherical, the structure of  $(\text{CH}_4)_3(\text{H}_2)_{25}$  is monoclinic with space group  $C2/m$  (coordinates are given in Supplemental




FIG. 2. (a) Representative x-ray diffraction patterns of the three compounds plotted as a function of exchanged wave vector and their Le Bail refinements. Refinements include  $(\text{CH}_4)_3(\text{H}_2)_{25}$ - $R\bar{3}m$  ( $a = 7.804 \text{ \AA}$  and  $c = 11.199 \text{ \AA}$ ),  $(\text{CH}_4)_2\text{H}_2$ - $I4/mcm$  ( $a = 7.195 \text{ \AA}$  and  $c = 5.909 \text{ \AA}$ ),  $\text{CH}_4$ - $A$ - $R3$  ( $a = 12.306 \text{ \AA}$  and  $c = 15.520 \text{ \AA}$ ),  $\text{CH}_4(\text{H}_2)_2$ - $P6_3/mmc$  ( $a = 4.981 \text{ \AA}$  and  $c = 8.125 \text{ \AA}$ ), and  $\text{Au}$ - $Fm\bar{3}m$  ( $a = 3.976 \text{ \AA}$ ). Excluded regions in the pattern of  $(\text{CH}_4)_3(\text{H}_2)_{25}$  correspond to rhenium (gasket material) and rhenium hydride (\*) and excess  $\text{H}_2$  (#). (b) Volumes per formula unit as a function of pressure. Symbols represent experimental data (the error bars are smaller than the symbol size), and black full lines represent their best second-order Birch-Murnaghan fits:  $\text{CH}_4(\text{H}_2)_{8.33}$ ,  $V_0 = 246 \pm 17 \text{ \AA}^3$  and  $K = 3.0 \pm 0.6 \text{ GPa}$ ;  $(\text{CH}_4)_2\text{H}_2$ ,  $V_0 = 101 \pm 2 \text{ \AA}^3$  and  $K = 10.2 \pm 0.8 \text{ GPa}$ ;  $\text{CH}_4(\text{H}_2)_2$ ,  $V_0 = 90 \pm 3 \text{ \AA}^3$  and  $K = 5.0 \pm 0.4 \text{ GPa}$ . Dashed lines represent volumes derived from our DFT calculations, and gray dotted lines represent the volumes of ideal mixtures of  $\text{CH}_4$ - $A$ / $\text{CH}_4$ - $B$  and  $\text{H}_2$ - $I$  using the previously determined equations of state [14,17,69]. The experimental volumes given in Ref. [21] are represented by gray circles. (c) Structural models of the three compounds, where  $\text{CH}_4$  and  $\text{H}_2$  are represented by brown and white spheres, respectively, and lines indicate  $\text{CH}_4$ - $\text{H}_2$  nearest neighbors.




FIG. 3. Representative vibrational Raman spectra of (a) 70%  $\text{H}_2$ , (b) 50%  $\text{H}_2$ , and (c) 30%  $\text{H}_2$  mixtures. Colors indicate the modes assigned to  $\text{CH}_4(\text{H}_2)_2$  (purple),  $(\text{CH}_4)_3(\text{H}_2)_{25}$  (green),  $(\text{CH}_4)_2\text{H}_2$  (orange), and excess  $\text{H}_2$  (black).






FIG. 4. Raman shift as a function of pressure of the  $\text{H}_2$  vibrational modes and C-H stretching modes of  $\text{CH}_4(\text{H}_2)_2$  and  $(\text{CH}_4)_3(\text{H}_2)_{25}$  (left panel, purple and green symbols, respectively), and  $(\text{CH}_4)_2\text{H}_2$  (right panel, orange symbols). Frequencies are collated from 30% (squares), 50% (circles), 70% (triangles), 80% (diamonds), and 90% (hexagons)  $\text{H}_2$  mixtures, with the most intense modes represented by larger symbols. Raman frequencies of individual mixtures are given in Supplemental Material [45]. The dot-dashed lines represent the calculated frequencies of the  $\text{H}_2$  modes and C-H stretching bands of each compound, with the thickest line representing the most intense modes. Black and gray solid lines represent pure  $\text{H}_2$  [6] and  $\text{CH}_4$  [15], respectively. The black dotted line represents the frequency of  $\text{H}_2$  in a Ne matrix [70]. The error in pressure ranges from  $\pm 0.2$  GPa below 50 GPa,  $\pm 1$  GPa below 150 GPa, and  $\pm 5$  GPa above. The error bars in frequency are smaller than the symbol size. Photomicrographs show examples of morphology of synthesised samples within the DAC sample chamber.

Material [45]). Le Bail refinements of the experimental diffraction patterns were performed using the trigonal space group  $R\bar{3}m$ , which is obtained considering the carbon atoms only, and the hexagonal setting, which resulted in good fits to the data. The obtained volume per methane molecule is close to  $V_{\text{CH}_4} + 8.33V_{\text{H}_2}$  (see Fig. 2), the volume of an ideal mixture of  $\text{CH}_4$  and  $\text{H}_2$  with the same composition. A comparison between the experimental and computational vibrational Raman modes of  $\text{CH}_4(\text{H}_2)_2$ - $P6_3/mmc$  and  $(\text{CH}_4)_3(\text{H}_2)_{25}$ - $C2/m$  is given in Fig. 4, where we see qualitatively good agreement between the number of modes, relative intensity, and frequency for both compounds.

In methane-rich concentrations of 30 mol %  $\text{H}_2$ , we observe a body-centered tetragonal  $\text{Al}_2\text{Cu}$ -type structure (space group  $I4/mcm$ ) to form above 4.6 GPa (with  $a = 7.195$  Å and  $c = 5.909$  Å at 5.4 GPa), which is consistent with the previously reported composition of  $(\text{CH}_4)_2\text{H}_2$  [21] (see Fig. 1). Raman spectroscopy measurements show that the  $\text{CH}_4$  stretching bands of  $(\text{CH}_4)_2\text{H}_2$  are close to the frequencies of pure solid  $\text{CH}_4$ , while two wagging modes are

observed. The number of modes and frequencies are in good agreement with our calculated values (see Fig. 4). The  $\nu_{1H}$  mode, corresponding to  $\text{H}_2$  units within the structure, exhibits the most extreme hardening of any stoichiometric molecular compound [25,28]. At a pressure of 207 GPa,  $(\text{CH}_4)_2\text{H}_2$ - $\nu_{1H}$  has a frequency of  $4798$   $\text{cm}^{-1}$ , over  $900$   $\text{cm}^{-1}$  higher than  $\text{H}_2$ - $\nu_1$  (see Fig. 4). Remarkably, the frequencies we observe reach similar values to an impurity  $\text{H}_2$  molecule isolated in a noble gas matrix, albeit at higher pressure [70]. Up to pressures of 207 GPa, we do not observe a turnover of  $\nu_{1H}$  (and 500 GPa in our calculations), unlike in pure  $\text{H}_2$ , which exhibits a maximum frequency at approximately 38 GPa [5].

These results highlight the enormous impact that the local environment around a  $\text{H}_2$  molecule can have on its vibron frequency. In Supplemental Material [45], we present a simple molecular model demonstrating that the vibron frequency of a  $\text{H}_2$  surrounded by two  $\text{CH}_4$  (respectively,  $\text{H}_2$ ) will continuously increase (respectively, decrease) if the surrounding molecules are pushed closer. A molecular orbital analysis reveals that, in  $\text{H}_2$ -dominated environments, intermolecular interactions lead to occupancy of antibonding H-H  $\sigma^*$  states, which does not happen in  $\text{CH}_4$ -dominated environments. In  $(\text{CH}_4)_2\text{H}_2$ , every  $\text{H}_2$  has two nearest  $\text{H}_2$  neighbors and eight nearest  $\text{CH}_4$  neighbors; as a result, the vibron exhibits extreme hardening [see Fig. 2(c) and Supplemental Material [45] for graphical representations of the unit cells and intermolecular distances]. This local environment is reversed in  $(\text{CH}_4)_3(\text{H}_2)_{25}$  with an average of 9.8  $\text{H}_2$  neighbors but only 2.24  $\text{CH}_4$  neighbors per  $\text{H}_2$  molecule; still, it remains very different from  $\text{H}_2$ -I, having a coordination number of 12 and considerably shorter  $\text{H}_2$ - $\text{H}_2$  distances [28]. For  $(\text{CH}_4)_3(\text{H}_2)_{25}$ , the  $\nu_{1H}$  mode has a maximum at about 70 GPa, and the vibron frequencies tend toward the values found for pure  $\text{H}_2$ . In  $\text{CH}_4(\text{H}_2)_2$ , where each  $\text{H}_2$  has six nearest  $\text{H}_2$  neighbors and six nearest  $\text{CH}_4$  neighbors, the vibron frequencies are interpolated between  $(\text{CH}_4)_3(\text{H}_2)_{25}$  and  $(\text{CH}_4)_2\text{H}_2$ .

We find that the pressure-composition phase diagram is simpler than previously reported [21], with only three stable  $\text{CH}_4$ - $\text{H}_2$  compositions. In 50 mol %  $\text{H}_2$  mixtures, we observe the coexistence of both  $(\text{CH}_4)_2\text{H}_2$  and  $\text{CH}_4(\text{H}_2)_2$ , together with some excess  $\text{H}_2$ , indicative that equilibrium has not been reached. Furthermore, we observe trace amounts of  $(\text{CH}_4)_3(\text{H}_2)_{25}$ , which can be attributed to macroscopic inhomogeneity within the sample chamber. In 70 mol % and 80 mol %  $\text{H}_2$  mixtures, we first observe the Raman signatures of  $(\text{CH}_4)_3(\text{H}_2)_{25}$  at 14.5 GPa and at 13.6 GPa (shown as open symbols in Fig. 1), respectively, while in 90 mol % mixtures, we observe  $(\text{CH}_4)_3(\text{H}_2)_{25}$  at 10.9 GPa. It is possible that, in 70 mol % and 80 mol %  $\text{H}_2$  mixtures, formation occurs at lower pressure but is below the detection limit in the Raman spectra. We find that even in 90 mol % mixtures the transformation from  $\text{CH}_4(\text{H}_2)_2$

and  $\text{H}_2$  to  $(\text{CH}_4)_3(\text{H}_2)_{25}$  is never complete, suggestive that the transformation is kinetically sluggish and due to hindered diffusion of  $\text{H}_2$  in the solid state. Although our liquidus curve is in good agreement with that reported previously, we do not observe either  $\text{CH}_4\text{H}_2$  and  $\text{CH}_4(\text{H}_2)_4$  and attribute this to a misinterpretation of  $\text{CH}_4(\text{H}_2)_2$  [21]. The experimental volume given for hexagonal wurtzite  $\text{CH}_4\text{H}_2$  is identical to our experimental volume for  $\text{CH}_4(\text{H}_2)_2$  at the same given pressure. Furthermore, after extensive structure searches, with randomly orientated molecules in the reported wurtzite center-of-mass positions, we do not find any version of the hexagonal wurtzite structure to be dynamically stable. The evidence for  $\text{CH}_4(\text{H}_2)_4$  was predominantly based on Raman spectroscopy, and we find to be a misinterpretation of the coexistence between the  $\text{CH}_4\text{-H}_2$  mixed fluid and solid  $\text{CH}_4(\text{H}_2)_2$  (see Fig. S8 [45]).

$(\text{CH}_4)_3(\text{H}_2)_{25}$  possesses 51.1 wt % molecular hydrogen (63.4 wt % if the hydrogen of methane is included), which is the highest hydrogen content of any currently known stoichiometric compound. Notwithstanding the good agreement between the experimental and calculated data,  $(\text{CH}_4)_3(\text{H}_2)_{25}$  represents an unusual and unique composition. All three compounds exhibit remarkable stability, with the Raman signatures of  $(\text{CH}_4)_2\text{H}_2$  and  $\text{CH}_4(\text{H}_2)_2$  detectable to at least 215 GPa and  $(\text{CH}_4)_3(\text{H}_2)_{25}$  up to 165 GPa. We estimate the melting temperatures  $T_m$  of  $(\text{CH}_4)_2\text{H}_2$  and  $\text{CH}_4(\text{H}_2)_2$  using Lindemann's equation based on calculated Debye temperatures and found  $T_m$  significantly higher than molecular hydrogen at all pressures (see Supplemental Material [45]). This would imply that hydrogen mixed with methane would potentially not possess the melting line turnover that is observed in pure hydrogen [7]. Taken together in a planetary context, this could influence critical properties of planetary matter such as thermal conductivities and viscosities. In a materials science context, high Debye temperatures coupled with extremely large vibron frequencies  $\nu_H$  are promising ingredients for high- $T_c$  superconductivity—provided a reasonable density of states  $N(E_F)$  at the Fermi energy. The molecular compounds described here are insulators (all remain transparent in the visible up to the highest pressures reached), but the presence of an electron or hole dopant would metallize the system, with partially charged entities  $\text{H}_2^{\pm\delta}$  present. This motif, at least in calculations, can lead to  $T_c$  close to 200 K [71].

Parts of this research were carried out at P02.2 at DESY, a member of the Helmholtz Association (HGF), and we thank H.-P. Liermann and K. Glazyrin for assistance. We also acknowledge the European Synchrotron Radiation Facility for provision of synchrotron radiation facilities at the ID15B beam line and assistance from M. Hanfland and D. Comboni. R. T. H. acknowledges that the project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 948895

“MetElOne”). M. P.-A. acknowledges the support of the UKRI Future leaders fellowship Mrc-Mr/T043733/1. L. J. C. was supported by the Engineering and Physical Sciences Research Council through the Condensed Matter Center for Doctoral Training (EP/L015110/1). Computational resources provided by the United Kingdom's National Supercomputer Service through the United Kingdom Car-Parrinello consortium (EP/P022561/1) and Project ID No. d56 “Planetary Interiors” and by the United Kingdom Materials and Molecular Modelling Hub (EP/P020194) are gratefully acknowledged. The authors thank J. Binns and F. A. Gorelli for useful discussions.

\*ross.howie@hpstar.ac.cn  
ross.howie@ed.ac.uk

- [1] W. B. Hubbard, W. J. Nellis, A. C. Mitchell, N. C. Holmes, S. S. Limaye, and P. C. McCandless, *Science* **253**, 648 (1991).
- [2] R. Chau, S. Hamel, and W. J. Nellis, *Nat. Commun.* **2**, 203 (2011).
- [3] S. S. Lobanov, P.-N. Chen, X.-J. Chen, C.-S. Zha, K. D. Litasov, H.-K. Mao, and A. F. Goncharov, *Nat. Commun.* **4**, 2446 (2013).
- [4] R. Helled and J. J. Fortney, *Phil. Trans. R. Soc. A* **378**, 20190474 (2020).
- [5] H.-k. Mao and R. J. Hemley, *Rev. Mod. Phys.* **66**, 671 (1994).
- [6] R. T. Howie, C. L. Guillaume, T. Scheler, A. F. Goncharov, and E. Gregoryanz, *Phys. Rev. Lett.* **108**, 125501 (2012).
- [7] R. T. Howie, P. Dalladay-Simpson, and E. Gregoryanz, *Nat. Mater.* **14**, 495 (2015).
- [8] P. Dalladay-Simpson, R. T. Howie, and E. Gregoryanz, *Nature (London)* **529**, 63 (2016).
- [9] E. Gregoryanz, C. Ji, P. Dalladay-Simpson, B. Li, R. T. Howie, and H.-K. Mao, *Matter Radiat. Extremes* **5**, 038101 (2020).
- [10] P. Loubeyre, F. Occelli, and P. Dumas, *Nature (London)* **577**, 631 (2020).
- [11] R. M. Hazen, H.-K. Mao, L. W. Finger, and P. M. Bell, *Appl. Phys. Lett.* **37**, 288 (1980).
- [12] H. Hirai, K. Konagai, T. Kawamura, Y. Yamamoto, and T. Yagi, *Phys. Earth Planet. Inter.* **174**, 242 (2009).
- [13] L. Sun, A. L. Ruoff, C.-S. Zha, and G. Stupian, *J. Phys. Chem. Solids* **67**, 2603 (2006).
- [14] L. Sun, W. Yi, L. Wang, J. Shu, S. Sinogeikin, Y. Meng, G. Shen, L. Bai, Y. Li, J. Liu, H.-K. Mao, and W. L. Mao, *Chem. Phys. Lett.* **473**, 72 (2009).
- [15] J. E. Proctor, H. E. Maynard-Casely, M. A. Hakeem, and D. Cantiah, *J. Raman Spectrosc.* **48**, 1777 (2017).
- [16] H. E. Maynard-Casely, C. L. Bull, M. Guthrie, I. Loa, M. I. McMahon, E. Gregoryanz, R. J. Nelmes, and J. S. Loveday, *J. Chem. Phys.* **133**, 064504 (2010).
- [17] H. E. Maynard-Casely, L. F. Lundegaard, I. Loa, M. I. McMahon, E. Gregoryanz, R. J. Nelmes, and J. S. Loveday, *J. Chem. Phys.* **141**, 234313 (2014).
- [18] L. J. Conway and A. Hermann, *Geosciences* **9**, 227 (2019).
- [19] M. Peña-Alvarez, A. V. Brovarone, M.-E. Donnelly, M. Wang, P. Dalladay-Simpson, R. Howie, and E. Gregoryanz, *Nat. Commun.* **12**, 6387 (2021).

- [20] P. Loubeyre, R. Letoullec, and J.-P. Pinceaux, *Phys. Rev. Lett.* **72**, 1360 (1994).
- [21] M. S. Somayazulu, L. W. Finger, R. J. Hemley, and H. K. Mao, *Science* **271**, 1400 (1996).
- [22] T. A. Strobel, P. Ganesh, M. Somayazulu, P. R. C. Kent, and R. J. Hemley, *Phys. Rev. Lett.* **107**, 255503 (2011).
- [23] M. Somayazulu, P. Dera, A. F. Goncharov, S. A. Gramsch, P. Liermann, W. Yang, Z. Liu, H.-K. Mao, and R. J. Hemley, *Nat. Chem.* **2**, 50 (2010).
- [24] D. K. Spaulding, G. Weck, P. Loubeyre, F. Datchi, P. Dumas, and M. Hanfland, *Nat. Commun.* **5**, 5739 (2014).
- [25] A. K. Kleppe, M. Amboage, and A. P. Jephcoat, *Sci. Rep.* **4**, 4989 (2014).
- [26] M. Somayazulu, P. Dera, J. Smith, and R. J. Hemley, *J. Chem. Phys.* **142**, 104503 (2015).
- [27] E. J. Pace, J. Binns, M. Peña Alvarez, P. Dalladay-Simpson, E. Gregoryanz, and R. T. Howie, *J. Chem. Phys.* **147**, 184303 (2017).
- [28] C. Ji, A. F. Goncharov, V. Shukla, N. K. Jena, D. Popov, B. Li, J. Wang, Y. Meng, V. B. Prakapenka, J. S. Smith, R. Ahuja, W. Yang, and H.-k. Mao, *Proc. Natl. Acad. Sci. U.S.A.* **114**, 3596 (2017).
- [29] D. Laniel, V. Svitlyk, G. Weck, and P. Loubeyre, *Phys. Chem. Chem. Phys.* **20**, 4050 (2018).
- [30] R. Turnbull, M.-E. Donnelly, M. Wang, M. Peña-Alvarez, C. Ji, P. Dalladay-Simpson, H.-k. Mao, E. Gregoryanz, and R. T. Howie, *Phys. Rev. Lett.* **121**, 195702 (2018).
- [31] J. Binns, P. Dalladay-Simpson, M. Wang, G. J. Ackland, E. Gregoryanz, and R. T. Howie, *Phys. Rev. B* **97**, 024111 (2018).
- [32] M. Ceppatelli, D. Scelta, M. Serrano-Ruiz, K. Dziubek, G. Garbarino, J. Jacobs, M. Mezouar, R. Bini, and M. Peruzzini, *Nat. Commun.* **11**, 6125 (2020).
- [33] W. L. Mao and H. K. Mao, *Proc. Natl. Acad. Sci. U.S.A.* **101**, 708 (2004).
- [34] W. L. Mao, V. V. Struzhkin, H. K. Mao, and R. J. Hemley, *Chem. Phys. Lett.* **402**, 66 (2005).
- [35] Y. Liu, D. Duan, F. Tian, X. Huang, D. Li, Z. Zhao, X. Sha, B. Chu, H. Zhang, B. Liu, and T. Cui, *RSC Adv.* **4**, 37569 (2014).
- [36] G. Saleh and A. R. Oganov, *Sci. Rep.* **6**, 32486 (2016).
- [37] A. S. Naumova, S. V. Lepeshkin, and A. R. Oganov, *J. Phys. Chem. C* **123**, 20497 (2019).
- [38] A. Zerr, G. Serghiou, R. Boehler, and M. Ross, *High Press. Res.* **26**, 23 (2006).
- [39] H. Kadobayashi, S. Ohnishi, H. Ohfuji, Y. Yamamoto, M. Muraoka, S. Yoshida, N. Hirao, S. Kawaguchi-Imada, and H. Hirai, *Sci. Rep.* **11**, 8165 (2021).
- [40] A. Kolesnikov, V. G. Kutcherov, and A. F. Goncharov, *Nat. Geosci.* **2**, 566 (2009).
- [41] D. A. Kudryavtsev, T. M. Fedotenko, E. G. Koemets, S. E. Khandarkhaeva, V. G. Kutcherov, and L. S. Dubrovinsky, *Sci. Rep.* **10**, 1483 (2020).
- [42] E. Snider, N. Dasenbrock-Gammon, R. McBride, M. Debessai, H. Vindana, K. Vencatasamy, K. V. Lawler, A. Salamat, and R. P. Dias, *Nature (London)* **586**, 373 (2020).
- [43] W. Cui, T. Bi, J. Shi, Y. Li, H. Liu, E. Zurek, and R. J. Hemley, *Phys. Rev. B* **101**, 134504 (2020).
- [44] A. F. Goncharov, E. Bykova, M. Bykov, X. Zhang, Y. Wang, S. Chariton, and V. B. Prakapenka, *J. Appl. Phys.* **131**, 025902 (2022).
- [45] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.128.215702> for further methodological details and additional experimental and computational figures, which includes Refs. [46–68].
- [46] E. Aprà *et al.*, *J. Chem. Phys.* **152**, 184102 (2020).
- [47] S. B. Ramsey, M. Pena-Alvarez, and G. J. Ackland, *Phys. Rev. B* **101**, 214306 (2020).
- [48] V. Labet, R. Hoffmann, and N. W. Ashcroft, *J. Chem. Phys.* **136**, 074502 (2012).
- [49] V. Labet, R. Hoffmann, and N. W. Ashcroft, *J. Chem. Phys.* **136**, 074503 (2012).
- [50] H.-K. Mao, J. Xu, and P. M. Bell, *J. Geophys. Res.* **91**, 4673 (1986).
- [51] Y. Akahama and H. Kawamura, *J. Appl. Phys.* **100**, 043516 (2006).
- [52] Y. Grin, F. R. Wagner, M. Armbrüster, M. Kohout, A. Leithe-Jasper, U. Schwarz, U. Wedig, and H. G. von Schnering, *J. Solid State Chem.* **179**, 1707 (2006).
- [53] S. Schaack, U. Ranieri, P. Depondt, R. Gaal, W. F. Kuhs, P. Gillet, F. Finocchi, and L. E. Bove, *Proc. Natl. Acad. Sci. U.S.A.* **116**, 16204 (2019).
- [54] H. T. Stokes, D. M. Hatch, and B. J. Campbell, FINDSYM, ISOTROPY Software Suite ([iso.byu.edu](http://iso.byu.edu)).
- [55] H. T. Stokes and D. M. Hatch, *J. Appl. Crystallogr.* **38**, 237 (2005).
- [56] T. Ohba, Y. Kitano, and Y. Komura, *Acta Cryst. C* **40**, 1 (1984).
- [57] K. Ishizaki, P. Bolsaitis, and I. Spain, *Solid State Commun.* **15**, 1591 (1974).
- [58] J. J. Gilvarry, *Phys. Rev.* **102**, 308 (1956).
- [59] F. A. Lindemann, *Phys. Z.* **11**, 609 (1910).
- [60] K. Refson, P. R. Tulip, and S. J. Clark, *Phys. Rev. B* **73**, 155114 (2006).
- [61] A. Tkatchenko and M. Scheffler, *Phys. Rev. Lett.* **102**, 073005 (2009).
- [62] G. Gao, A. R. Oganov, Y. Ma, H. Wang, P. Li, Y. Li, T. Iitaka, and G. Zou, *J. Chem. Phys.* **133**, 144508 (2010).
- [63] S. J. Clark, M. D. Segall, C. J. Pickard, P. J. Hasnip, M. I. J. Probert, K. Refson, and M. C. Payne, *Z. Kristallogr.* **220**, 567 (2005).
- [64] K. Momma and F. Izumi, *J. Appl. Crystallogr.* **44**, 1272 (2011).
- [65] J. Gonzalez-Platas, M. Alvaro, F. Nestola, and R. Angel, *J. Appl. Crystallogr.* **49**, 1377 (2016).
- [66] J. Rodríguez-Carvajal, *Physica (Amsterdam)* **192B**, 55 (1993).
- [67] Y. Fei, A. Ricolleau, M. Frank, K. Mibe, G. Shen, and V. Prakapenka, *Proc. Natl. Acad. Sci. U.S.A.* **104**, 9182 (2007).
- [68] C. Prescher and V. B. Prakapenka, *High Press. Res.* **35**, 223 (2015).
- [69] P. Loubeyre, R. LeToullec, D. Hausermann, M. Hanfland, R. J. Hemley, H. K. Mao, and L. W. Finger, *Nature (London)* **383**, 702 (1996).
- [70] P. Loubeyre, R. Letoullec, and J. P. Pinceaux, *Phys. Rev. Lett.* **67**, 3271 (1991).
- [71] B. Chen, L. J. Conway, W. Sun, X. Kuang, C. Lu, and A. Hermann, *Phys. Rev. B* **103**, 035131 (2021).