



# **An inverted textbook on thermodynamics: Part I**

Graeme Ackland

GRAEME ACKLAND

---

# **AN INVERTED TEXTBOOK ON THERMODYNAMICS: PART I**

An inverted textbook on thermodynamics: Part I

2<sup>nd</sup> edition

© 2019 Graeme Ackland & [bookboon.com](http://bookboon.com)

ISBN 978-87-403-3170-7

# CONTENTS

|           |                                                                |           |
|-----------|----------------------------------------------------------------|-----------|
| <b>1</b>  | <b>Questions: Some properties of materials</b>                 | <b>6</b>  |
| <b>2</b>  | <b>Questions: Temperature scales, work, equations of state</b> | <b>9</b>  |
| <b>3</b>  | <b>Questions: Work and heat, the First Law</b>                 | <b>11</b> |
| <b>4</b>  | <b>Questions: Cycles and the Second Law</b>                    | <b>14</b> |
| <b>5</b>  | <b>Questions: Thermodynamic Potentials</b>                     | <b>16</b> |
| <b>6</b>  | <b>Questions: Thermodynamic Potentials</b>                     | <b>17</b> |
| <b>7</b>  | <b>Questions: Expansion Processes</b>                          | <b>19</b> |
| <b>8</b>  | <b>Questions: Thermodynamics in other systems</b>              | <b>21</b> |
| <b>9</b>  | <b>Questions: Phase transitions</b>                            | <b>24</b> |
| <b>10</b> | <b>Questions: Chemical Potential</b>                           | <b>27</b> |

|           |                                                     |                |
|-----------|-----------------------------------------------------|----------------|
| <b>11</b> | <b>Some properties of materials</b>                 | <b>Part II</b> |
| <b>12</b> | <b>Temperature scales, work, equations of state</b> | <b>Part II</b> |
| <b>13</b> | <b>Work and heat, the First Law</b>                 | <b>Part II</b> |
| <b>14</b> | <b>Cycles and the Second Law</b>                    | <b>Part II</b> |
| <b>15</b> | <b>Entropy</b>                                      | <b>Part II</b> |
| <b>16</b> | <b>Thermodynamic Potentials</b>                     | <b>Part II</b> |
| <b>17</b> | <b>Expansion Processes</b>                          | <b>Part II</b> |
| <b>18</b> | <b>Thermodynamics in other systems</b>              | <b>Part II</b> |
| <b>19</b> | <b>Phase transitions</b>                            | <b>Part II</b> |
| <b>20</b> | <b>Chemical Potential</b>                           | <b>Part II</b> |

# 1 Questions: Some properties of materials

Use the following values where necessary in the questions:

|                                        |   |                                          |
|----------------------------------------|---|------------------------------------------|
| latent heat of fusion (melting) of ice | = | 334 kJ kg <sup>-1</sup>                  |
| latent heat of vaporization of water   | = | 2256 kJ kg <sup>-1</sup>                 |
| specific heat capacity of ice          | = | 1.94 kJ kg <sup>-1</sup> K <sup>-1</sup> |
| specific heat capacity of water        | = | 4.2 kJ kg <sup>-1</sup> K <sup>-1</sup>  |
| specific heat capacity of steam        | = | 2.04 kJ kg <sup>-1</sup> K <sup>-1</sup> |

## 1. Orders of Magnitude Place the following quantities of energy in order.

- 1/ The daily energy consumption of the UK
- 2/ The binding energy of all the electrons in one kg of hydrogen molecules.
- 3/ The rest mass energy of one kg of hydrogen molecules.
- 4/ The zero point energy of one kg of hydrogen molecules (vibrational frequency 4161cm<sup>-1</sup>).
- 5/ The energy released when one kg of deuterium molecules fuse to create one kg of helium.
- 6/ The thermal energy of one kg of hydrogen molecules at 300K.
- 7/ The calorific value of a cold chicken sandwich at 280K.
- 8/ The additional thermal energy of a cold chicken sandwich at 320K.
- 9/ The energy required to remove the chicken sandwich from the earth's gravitational field.
- 10/ The kinetic energy of the chicken sandwich on a train at 50m/s.
- 11/ 1kg of coal, when burnt.
- 12/ 1kg of uranium-235, when fissioned
- 13/ 1000 cubic metres of air moving at 10m/s
- 14/ 1000 cubic metres of water, raised by 5m

## 2. Heating and metabolism

A class of 180 students sits in a lecture theatre for one hour, each student metabolising at 100 W. The lecture theatre is a cubic room twenty metres long on each side and 2.5 metres high. The specific heat capacity of air at constant volume is 1 kJ kg<sup>-1</sup> K<sup>-1</sup>. The density of air is about 1.2 kg m<sup>-3</sup>. The initial temperature was 20°C. The ventilation is poor, the air conditioning is broken and the walls are well insulated. What is the room temperature at the end of the lecture?

Compare this to the rate of heat production of the sun, which produces  $3.86 \times 10^{26}$ W.

## 3. Thermal properties in food

A 500-gram box of strawberries is cooled in a refrigerator, from an initial temperature of 25 °C down to the fridge temperature of 4 °C.

- (a) *Estimate* how much heat is removed from the strawberries during the cooling, explaining your reasoning. *Hint: Strawberries have a water content of about 88%. For this and subsequent problems, some of the data given at the start of this section may be useful.*
- (b) Fruits and vegetables actually respire continuously whilst in storage, taking in oxygen and converting it to carbon dioxide. In the case of strawberries take the heat produced by this reaction to be about 210 mW kg<sup>-1</sup>. How does your estimate of heat removed change if respiration is taken into account? *Hint: You need to make a sensible assumption about timescales.*
- (c) The strawberries are now removed from the fridge and put into a polystyrene (i.e. thermally insulating) container in the kitchen. How long will it take for the strawberries to reach room temperature?
- (d) The nutritional value of 100g of strawberries is 33kcal (140kJ). Use this and the respiration rate to estimate the lifetime of strawberries.

4. **Phase changes: latent heat.**

An ice cube of mass 0.03 kg at 0°C is added to 0.2 kg of water at 20°C in an insulated container.

(a) Does all the ice melt? (b) What is the final temperature of the drink?

*Comment: The ice does all melt, but specify carefully the criterion which must be satisfied.*

5. **Conservation of energy: gravity and heat**

In 1845 James Prescott Joule suggested that the water at the bottom of a waterfall should be warmer than at the top. In particular, for Niagara Falls (a height of about 50 m) the temperature difference would be approximately 0.12 °C.

How would you go about calculating this number?

How do you know this must be an overestimate?

Unlike in 1845 most of the water nowadays is diverted through hydroelectric power schemes. How does this affect Joule's prediction?

6. **The ideal gas law**

The ideal gas is defined by its *equation of state*, which relates the variables pressure  $P$ , temperature  $T$  and volume  $V$ :

$$PV = nRT$$

Here  $n$  is the number of moles of the gas sample, and  $R$  is the gas constant. Calculate the volume occupied by a sample of ideal gas at atmospheric pressure and a temperature of 25°C, given that its volume under *standard temperature and pressure* is  $V_m = 2.2414 \times 10^{-2} \text{ m}^3$ .

n.b. Standard temperature and pressure means  $T = 0^\circ\text{C}$ ,  $P = 1 \text{ atm} = 101325 \text{ Pa}$ .

7. **Another 'ideal' gas** You are told that at a pressure of 1.2 atm and temperature of  $T = 300 \text{ K}$ ,  $n$  moles of oxygen occupy a volume of  $82 \text{ cm}^3$ . Calculate  $n$  and the *mass* of the oxygen sample, assuming that under these conditions the gas behaves ideally. How would the answer change if the sample was *ozone* ( $\text{O}_3$ )?

8. **Melting, heating and boiling.** A 1 kg sample of ice at an initial temperature of  $-4^\circ\text{C}$  is heated at constant pressure in an insulated container, heat being supplied at a constant rate of  $1 \text{ kJ s}^{-1}$ , until the sample is steam at  $110^\circ\text{C}$ . Using the data given below, sketch the temperature as a function of time. Discuss the differences in the times taken for melting, heating, and boiling.

9. **Conduction of heat (an idealised hot water bottle).** A certain heat source (a *heat reservoir* or *thermal reservoir*), is always at temperature  $T_0$ . Heat is transferred from it through a slab of thickness  $L$  to an object which is initially at a temperature  $T_1 < T_0$ . The object, which is otherwise thermally insulated from its surroundings, has a mass  $m = 0.5 \text{ kg}$  and a specific heat capacity  $c = 4 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$ . Heat is conducted through the slab at a rate (in  $\text{J s}^{-1}$ ) specified by the formula  $KA((T_0 - T)/L)$ , where  $K$  is the thermal conductivity of the slab,  $A$  is the area of the slab through which heat is transferred and  $T$  is the instantaneous temperature of the object.

(a) Show that provided certain assumptions are made,

$$KA((T_0 - T)/L)\Delta t = mc\Delta T$$

where the temperature of the object changes from  $T$  to  $T + \Delta T$  during the time interval  $\Delta t$ . (For simplicity, treat the arrangement as a 1-dimensional system with the  $x$ -axis perpendicular to the slab.)

(b) Show that for small  $\Delta t$ , and hence small  $\Delta T$ , the above equation can be rearranged and integrated to give

$$T_2 - T_1 = (T_0 - T_1) \left( 1 - \exp \left( -\frac{KA(t_2 - t_1)}{Lmc} \right) \right)$$

$T_2$  and  $T_1$  are the temperatures of the object at the end,  $t = t_2$ , and the beginning,  $t = t_1$ , respectively, of the process described.

- (c) Put  $A = 100 \text{ cm}^2$  and  $L = 1 \text{ cm}$ . Calculate the value of  $T_2 - T_1$  for  $t_2 - t_1$  equal to (a) 2 s and (b) 2000 s (about half-an-hour) for  $T_0 = 60^\circ\text{C}$  and  $T_1 = 35^\circ\text{C}$  for the following slab materials: whose thermal conductivities are given in the table.

aluminium (  $K = 200 \text{ W m}^{-1} \text{ K}^{-1}$  ) porcelain (  $K = 1.5 \text{ W m}^{-1} \text{ K}^{-1}$  ) rubber (  $K = 0.15 \text{ W m}^{-1} \text{ K}^{-1}$  ) wool (  $K = 0.05 \text{ W m}^{-1} \text{ K}^{-1}$  ) air (  $K = 0.025 \text{ W m}^{-1} \text{ K}^{-1}$  )

*Comment: The mathematical analysis incorporates a step which is quite common in problems in thermodynamics. If the temperature of part of a composite system changes from  $T_i$  to  $T_f$ , work done and/or heat flow during the process can often be calculated most quickly by considering intermediate stages in which the temperature changes from  $T$  to  $T + dT$ , setting up the appropriate equations and integrating them.*

## 2 Questions: Temperature scales, work, equations of state

### 1. Work done in other processes

- In melting:** Ice at  $0^\circ\text{C}$  and at a pressure 1 atm, has a density of  $916\text{ kg m}^{-3}$ , while water under these conditions has a density  $1000\text{ kg m}^{-3}$ . How much work is done when 10 kg of ice melts into water? Explain why this work is done “by the atmosphere” rather than “against the atmosphere”. Is there a difference between latent heat of melting at constant pressure, and latent heat of melting at constant volume?
- On a wire:** Calculate the work done when a copper wire of length 10cm holding a 2kg weight extends by 0.1% due to reversible heating. Given the linear thermal expansion coefficient is  $16.6 \times 10^{-6}\text{ K}^{-1}$ , estimate the required temperature change.
- In a wire:** Calculate the electrical work done when the wire is connected to a 6V battery for 10sec. (assume resistance=4.2m $\Omega$ )

### 2. Calculating properties from the equation of state, and vice versa

- The isothermal compressibility  $\kappa$  and the isobaric volume expansivity  $\beta$  are given by:

$$\kappa = -\frac{1}{V} \left( \frac{\partial V}{\partial P} \right)_T \quad \beta = \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_P$$

Using the equation of state, calculate the isobaric expansivity,  $\beta$  and isothermal compressibility,  $\kappa$ , of an ideal gas. Show that the bulk modulus for an ideal gas,  $K = P$

- A substance is found to have an isothermal bulk modulus  $K = v/a$  and an isobaric expansivity  $\beta = 2bT/v$  where  $a$  and  $b$  are constants and  $v$  is the molar volume. Show that the equation of state is  $v - bT^2 + aP = \text{a constant}$ .

[Hint: Integrate the expressions involving partial derivatives for  $\kappa$  and  $\beta$ , and merge/reconcile the outcomes.]

- Show that for the ideal gas, the difference between the heat capacities at constant volume and constant pressure is actually given by:

$$C_P - C_V = \frac{VT\beta^2}{\kappa}$$

Remember  $C_P - C_V = R$  and  $PV = RT$  for one mole of an ideal gas.

### 3. Reversible adiabatic expansion of ideal gas

(This question involves working through the final section of lecture 3)

Explain why the first Law for an reversible adiabatic process gives  $\Delta U = -PdV$ , and why this equation doesn't hold for the Joule expansion.

Assuming that for an ideal gas  $U = C_V T$ , prove that the First Law leads to the statement that  $PV^\gamma$  is constant in a reversible adiabatic process.

A container of Helium and a container of Oxygen are initially the same pressure and temperature. Each undergoes a reversible adiabatic compression to half its original volume. Assuming they are ideal gases, how do their temperatures compare?

### 4. Temperature scales: based on water

At atmospheric pressure, water has a density of  $960\text{ kg/m}^3$  at  $100^\circ\text{C}$ , the same as supercooled water at  $-40^\circ\text{C}$ . The maximum density,  $1000\text{ kg/m}^3$ , is at  $4^\circ\text{C}$ .

If the equation approximating an isobar is  $\rho = A + BT + CT^2 + DT^3$ , evaluate the constants A-D. (Hint - you are free to choose the zero of the temperature scale as you please)

A water thermometer comprises water in a glass tube of constant radius. At  $4^\circ\text{C}$ , the column of water is 100mm high. At what temperature will it be 101mm high?

Give reasons why mercury is used in preference to water in thermometers.

**5. Joule's experiment**

In an experiment similar to Joule's paddle wheel experiment, a mass of 20 kg drops slowly through a distance of 2 m, driving the paddles immersed in 2 kg of water. Viscous dissipation generates heat in the water.

- (a) Ignoring heat losses, bearing friction, etc, calculate the rise in temperature of the water.
- (b) What would be the error in determining the mechanical equivalent of heat if the mass was still moving at  $10 \text{ cm s}^{-1}$  when it hits the ground?

(The heat capacity of water is  $4.2 \text{ kJ K}^{-1}\text{kg}^{-1}$  )

**6. Temperature scale based on electrical resistance**

Idiosyncratic Roger invents a temperature scale using as his thermometric property the resistance  $R(T)$  of a special wire. He decides to calibrate his scale using the ice temperature (273.15K) and the triple point of water (273.16K). His wire has resistance  $R_0$  at the ice point temperature, which he defines as  $T_R = 273.15$ . It has resistance  $R_0 + \Delta R$  at the triple point, which he defines as  $T_R = 273.16$ . He then defines other temperatures by  $T_R = 273.15 + 0.01(R - R_0)/\Delta R$ , which he determines by measuring the resistance  $R$ .

Unbeknownst to Roger, the resistance of his wire is given by

$$R = R_0(1 + \alpha T + \beta T^2)$$

where  $T$  is the temperature in degrees Celsius measured on the ideal gas scale. The constants  $\alpha$  and  $\beta$  are  $3.8 \times 10^{-3} \text{ K}^{-1}$  and  $-3.0 \times 10^{-6} \text{ K}^{-2}$  respectively. What temperature on Roger's resistance scale corresponds to a temperature of  $70^\circ\text{C}$  on the ideal gas scale?

**7. Temperature scales: influence of thermal properties**

An alcohol and a mercury thermometer are constructed so that they agree at temperatures of  $0^\circ\text{C}$  and  $100^\circ\text{C}$ , and each scale (i.e. column of alcohol or mercury) is marked with 100 equal divisions between these two 'fixed points'. Will the two scales necessarily give the same reading at all temperatures between the fixed points? Explain your reasoning. What conditions are necessary for the two scales to agree completely? What conditions are needed for the thermometers to agree and *also* give readings in Celsius.

### 3 Questions: Work and heat, the First Law

#### 1. Work done in the expansion of an ideal gas


The volume of a given system containing  $n$  moles of a monatomic ideal gas is given by  $V = V(P, T)$ , where  $P$  is the pressure and  $T$  the temperature, and  $PV = nRT$ .

- (a) Obtain an expression in terms of the change in pressure,  $dP$ , for the incremental work done on the system,  $dW$  in an isothermal expansion.
- (b) The gas expands isothermally to twice its original volume. Using the formula for the incremental work on the system in terms of  $dP$  (previous part), obtain an expression for the total work done on the system.
- (c) Write down an expression in terms of the change of temperature,  $\Delta T$ , for the work done on the system in an *isobaric* expansion.

#### 2. Work, heat, $PV$ diagrams

A gas, contained in a cylinder fitted with a frictionless piston, is taken from the state A to the state B along the path ACB shown in the diagram. On this path, 80 J of heat flows into the system and the system **does** 30 J of work.

- Using the First Law, write down the difference in internal energy between the states A and B.
- How much heat flows into the system for the process represented by the path ADB if the work done by the system on this path is 10 J?
- When the system returns from state B to state A along the **curved** path AB, the work done **on** the system is 20 J. What is the heat transfer?
- If the internal energy of the system in state A is  $U_A$  while in the state D,  $U_D = U_A + 40$  J, find the heat absorbed in the processes AD and DB.



### 3. Free expansion of a van der Waals gas

Assuming that helium obeys the van der Waals equation of state, determine the change in temperature when one kilomole of helium gas, initially at 20°C and with a volume of 0.12 m<sup>3</sup>, undergoes a free expansion to a final pressure of one atmosphere. You should use the following expression:

$$\left(\frac{\partial T}{\partial V}\right)_U = -\frac{a}{C_V} \left(\frac{n}{V}\right)^2$$

For this gas the relevant parameters in the van der Waals equation of state are given by  $a = 3.44 \times 10^3$  J m<sup>3</sup> kmol<sup>-2</sup>;  $b = 0.0234$  m<sup>3</sup> kmol<sup>-1</sup>; and in this case  $C_V/nR = 1.506$ .

(Hint: You may approximate. First show that the initial pressure is much higher than the final one. Then you may assume that the final volume is much larger than the initial volume.)

Numerical answer: -2.3 K.

### 4. Free expansion experiments and internal energy of ideal gas

In experiments on the free expansion of low-pressure (i.e. 'ideal') gases it was found that there was no measurable temperature change of the gas, i.e.

$$\left(\frac{\partial T}{\partial V}\right)_{U,ideal} = 0$$

Show that this observation means that the internal energy of the gas,  $U$ , must be a function of temperature  $T$  only. Hint: you will need to use the reciprocity relation between partial derivatives:

$$\left(\frac{\partial x}{\partial y}\right)_z \left(\frac{\partial y}{\partial z}\right)_x \left(\frac{\partial z}{\partial x}\right)_y = -1$$

In this case the relevant variables (x,y,z) are  $T$ ,  $U$ , and  $V$ .

### 5. Adiabatic processes

Use the First Law and the Ideal Gas equation to show that the pressure and volume of an ideal gas in a reversible adiabatic expansion are related by  $PV^\gamma = c$  where  $c$  and  $\gamma = C_P/C_V$  are constants. Show that the work done by the gas in a reversible adiabatic expansion from  $(P_1, V_1)$  to  $(P_2, V_2)$  is

$$W = \frac{P_1 V_1 - P_2 V_2}{\gamma - 1}$$

Hint: Remember that  $c_P - c_V = R$  for 1 mole of an ideal gas, and use the Ideal Gas equation in differential form  $RdT = PdV + VdP$ .

### 6. Isobaric processes

Use the First Law to find the change in internal energy of a monatomic ideal gas in an isobaric expansion at 1 atm from a volume of 5 m<sup>3</sup> to a volume of 10 m<sup>3</sup>. Show that this is independent of the number of moles of gas.  $\gamma$  for a monatomic ideal gas is 5/3.

**7. Cooling in expansions**

By considering adiabats and isenthalps on an indicator diagram, *explain* why the adiabatic expansion produces more cooling than either a free expansion (Joule process) or a throttling (Joule-Kelvin) process for an ideal gas or similar material.

Given that:

$$\left(\frac{\partial H}{\partial P}\right)_T = V - T \left(\frac{\partial V}{\partial T}\right)_P$$

use the triple product rule for partial differentials to derive the expression

$$\mu_{JK} = \left(\frac{\partial T}{\partial P}\right)_H = \frac{1}{C_P} \left( T \left(\frac{\partial V}{\partial T}\right)_P - V \right)$$

Show that over a range of  $T$  where  $\mu_{JK}$  is independent of temperature, the cooling in a throttling (Joule-Kelvin) process with a pressure change from  $P_1$  to  $P_2$  is

$$\Delta T = \int_{P_1}^{P_2} \left(\frac{\partial T}{\partial P}\right)_H dP$$

In a similar way, show that the cooling in an adiabatic reversible expansion from a pressure  $P_1$  to  $P_2$  is

$$\Delta T = \int_{P_1}^{P_2} \left(\frac{\partial T}{\partial P}\right)_S dP$$

Hence *prove* that, for a given pressure change, the adiabatic expansion produces more cooling than a throttling process, i.e. that the difference in the integrands  $(\partial T/\partial P)_S - (\partial T/\partial P)_H$  is positive.)

## 4 Questions: Cycles and the Second Law

### 1. Statements of the Second Law of Thermodynamics

Show that if the Clausius statement of the Second Law of Thermodynamics is false, the Kelvin-Planck statement of the Second Law of Thermodynamics must be false also.


*Hint: Make a composite heat engine consisting of a Clausius-statement-breaking refrigerator which transfers an amount of heat  $Q_2$  per cycle from a cold to a hot body together with an engine which delivers the same amount of heat,  $Q_2$ , per cycle to the cold body.*

### 2. Efficiency of engines part 1

Which gives the greater increase in the efficiency of a Carnot engine: increasing the temperature of the hot reservoir or lowering the temperature of the cold reservoir by the same amount?

### 3. Efficiency of engines part 2

The maximum efficiency of a heat engine can be increased by reducing the temperature of the lower temperature reservoir. Consider the composite engine shown in the diagram opposite. Engine E operates between a high temperature  $T_1$  and a body at  $T_2$ , lower than the ambient temperature. This lower temperature body is maintained at lower than ambient temperature by a refrigerator which extracts heat  $Q_2$  from the lower temperature body and emits heat at the ambient temperature  $T_a$ . Show that the composite engine ER has a maximum efficiency equal to that of a single engine operating between the temperatures  $T_1$  and  $T_a$ , in other words that nothing is gained by artificially cooling the lower temperature reservoir.




For the composite engine use the usual formula for efficiency, in terms of the **net** work done on the surroundings and the (heat) energy supplied at the **highest** temperature. Then exploit the relationships between heat flows and temperatures.

### 4. Efficiency of Engines part 3

An ideal gas is taken through the reversible cycle (The Otto cycle) shown in the diagram where **ab** and **cd** are adiabatics. The temperature at *a* is  $T_a$  and so on.

Briefly describe the working cycle identifying the processes during which heat flows in or out of the system. Calculate expressions for these heat flows in terms of an appropriate heat capacity. Give an expression for the net work output per cycle and identify this quantity on a sketch of the  $P - V$  diagram.



The 'Otto cycle' is an idealisation of the internal combustion engine. Which segments represent fuel compression, burning, and exhaust?

Show that the efficiency of the Otto cycle is

$$1 - \frac{T_d - T_a}{T_c - T_b}$$

5. **Efficiency of Engines part 4** An engineer applies to a venture capital company for money to market a new heat engine, which is claimed to extract 5000 J of heat from a reservoir at 400 K, reject 3500 J to a reservoir at 300 K, and do 1500 J of work per cycle on the surroundings. How should the venture capitalists go about studying details of the device.
6. **The best possible fridge** Using reasonable values for the temperatures inside and outside a domestic refrigerator, calculate its maximum possible efficiency.

### 7. Domestic heating

A building is heated to  $27^\circ\text{C}$ . How much heat per second could be supplied by

- (a) An electric heater, with power input 20kW
- (b) A heat pump connected to an adjacent river at  $7^\circ\text{C}$ , with power input 20kW.

### 8. Multipurpose device.

A company markets a device which, it claims, can extract 400W heat from a fridge compartment *and* deliver 1kW heating to the living room using just 100W of electricity. Consider both First and Second Laws to determine whether the claim is thermodynamically plausible?

### 9. Yet another cycle

An engine cycle using an ideal gas consists of the following steps

- (i) an isobaric compression from Volume  $V_a$  to  $V_b$  at pressure  $P_a$
- (ii) an increase in pressure from  $P_a$  to  $P_b$  at a constant volume  $V_b$
- (iii) an adiabatic expansion from  $(P_b, V_b)$  to the original state at  $(P_a, V_a)$

Sketch this cycle on a PV plot. Describe the steps where heat enters and leaves the system.

What is the efficiency of a heat engine expressed in terms of the magnitudes of the heat inputs and outputs? Show that the efficiency of the cycle described above is

$$\eta = 1 - \frac{\gamma P_a}{V_b} \frac{V_a - V_b}{P_b - P_a}$$


why is it peculiar to see  $\gamma$  in this expression?

### 10. Work extracted while approaching thermal equilibrium

A small Carnot engine operates between two identical bodies each having a finite heat capacity  $C_P$ , initially at temperatures  $T_1$  and  $T_2$  respectively, as indicated in the diagram.

Heat flows from the higher temperature body to the lower temperature body until the two bodies eventually reach the same temperature,  $T_f$ . Calculate the total amount of work done by the Carnot engine before the temperatures of both bodies reach  $T_f$ .

How would the result differ if the cold reservoir was large enough that its temperature remains constant?



*Hint: It is necessary to specify an intermediate stage between start (here the two temperatures  $T_1$  and  $T_2$ ) and finish (here the common temperature  $T_f$ ). To avoid an ambiguous notation, use  $T'$  as the intermediate temperature for the body whose initial temperature was  $T_1$ ; and  $T''$  as the intermediate temperature for the body whose initial temperature was  $T_2$ . Then small temperature changes for an intermediate cycle can be represented – loosely – by  $dT'$  and  $dT''$ , and heat flows by  $C_P dT'$  and  $C_P dT''$ . However, great care has to be taken when allocating/associating signs with these four quantities! An application of the general relationship between heat flows and temperatures, in this context, provides an equation which can be integrated between limits denoted by  $T_1$ ,  $T_2$  and  $T_f$ , as appropriate. The relationship between  $T_f$ ,  $T_1$  and  $T_2$  turns out to be  $T_f^2 = T_1 T_2$ , which you should derive.*

## 5 Questions: Thermodynamic Potentials

### 1. Calculation of entropy change

5 kg of hot water at 25°C cools to the 5°C temperature of its surroundings. How much heat flows? What is the entropy change of the surroundings? What is the entropy change of the water? ( $c_P$  for water =  $4.19 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$ .)

### 2. Variation on the same theme

An electric current of 10 A flows through a resistor of 20 ohms which is kept at 27°C by being immersed in running water. What is the entropy change per second of the resistor, the water and the universe?

### 3. Entropy and ideal gases

The internal energy of an ideal gas is all kinetic energy, and so it depends only on the temperature.

Use this fact to derive an expression for the difference,  $S_2 - S_1$ , in the entropies  $S_1$  and  $S_2$  of one mole of an ideal gas at volumes  $V_1$  and  $V_2$  respectively, at the same temperature.

Then show that for an ideal gas,  $C_P$  is independent of pressure  $P$ .

### 4. General entropy changes in a ideal gas

The heat capacity of an ideal gas is not fully defined by the equation of state, it depends on details of the molecule. Consider the case of a near ideal gas with equation of state  $PV = nRT$  and  $c_v = A + BT$  where  $A$  and  $B$  are constants, and show that the change in entropy between state  $(V_1, T_1)$  and state  $(V_2, T_2)$  is

$$\Delta S = A \ln(T_2/T_1) + B(T_2 - T_1) + R \ln(V_2/V_1)$$

*Hint: From the Central Equation,  $dS$  can be written as a sum of two terms, one involving  $dU$  and the other  $dV$ . A glance at the result sought shows that the term involving  $dU$  has to be re-cast as one involving  $dT$  before doing the integration to get  $\Delta S$ .*

### 5. Another look at the Carnot cycle

Sketch a Carnot cycle, not on a  $PV$  diagram but on a  $TS$  diagram (i.e. temperature versus entropy). Show that the area within the closed path is equal to the heat absorbed per cycle provided that the path is traced out clockwise.

*Hint: consider what is happening on each 'leg' of the Carnot cycle, in terms of entropy and temperature.*

### 6. Entropy change in a (simplified) gin and tonic

A cube of ice of mass 30 g melts in a glass of water at 0°C in a room at 20°C. The water is stirred slowly to keep its temperature at 0°C, but gently enough that the work done can be neglected. Calculate the changes in entropy of the ice, the water and the air in the room. (The latent heat of fusion of ice is  $334 \text{ kJ kg}^{-1}$ .)

### 7. Entropy change, reversible and irreversible processes

A block of lead with heat capacity  $C_P = 1000 \text{ J K}^{-1}$  is cooled from  $T_1 = 200 \text{ K}$  to  $T_2 = 100 \text{ K}$  by:

- plunging the block into a large bath of liquid at 100 K,
- first plunging the block into a large bath at 150 K, equilibrating, then plunging into a second bath at 100 K.
- a reversible process

Calculate the entropy change of the universe in each case, and give an explanation for how the reversible process could be implemented in practice.

## 6 Questions: Thermodynamic Potentials

### 1. Heat Capacities

The heat capacity is *defined* as the amount of heat required to raise the temperature of a body by 1K. From this, the First Law and the Central Equation, prove that:

$$C_V = \left( \frac{\partial U}{\partial T} \right)_V = T \left( \frac{\partial S}{\partial T} \right)_V$$

$$C_P = \left( \frac{\partial H}{\partial T} \right)_P = T \left( \frac{\partial S}{\partial T} \right)_P$$

### 2. Helmholtz function and pressure

Write down the differential form of the Helmholtz function  $F = U - TS$ , and an expression for pressure in terms of  $F$ .

The specific Helmholtz function of a particular gas is:

$$f = \frac{F}{n} = f_0(T) - \frac{a}{v} - RT \ln(v - b)$$

where  $f_0$  is a function of  $T$  only, while  $a$  and  $b$  are constants. Calculate the pressure of the gas, and hence its equation of state.

### 3. Using Maxwell relations

The Helmholtz thermodynamic potential  $F$  is defined by  $F = U - TS$ .

Starting from the Central equation ( $dU = -PdV + TdS$ ) and this definition, derive the Maxwell relation associated with the Helmholtz potential.

For any material, the difference between the heat capacities at constant pressure and volume is given by

$$C_P - C_V = \left[ \left( \frac{\partial U}{\partial V} \right)_T + P \right] \left( \frac{\partial V}{\partial T} \right)_P$$

Using the Maxwell relation, show that

$$\left( \frac{\partial C_V}{\partial V} \right)_T = T \left( \frac{\partial^2 P}{\partial T^2} \right)_V$$

and that

$$C_P - C_V = \frac{VT\beta_P^2}{\kappa_T}$$

where  $\beta_P$  is the isobaric volume expansivity and  $\kappa_T$  is the isothermal compressibility.

Use a similar technique to prove that the difference between the isothermal and the adiabatic compressibilities is

$$\kappa_T - \kappa_S = \frac{TV\beta_P^2}{C_P}$$

Verify that  $C_P - C_V = R$  and  $\kappa_T - \kappa_S$  for a monatomic ideal gas.

### 4. A block of metal.

A block of metal is subjected to an adiabatic and reversible increase of pressure from  $P_1$  to  $P_2$ . Show that the initial and final temperatures  $T_1$  and  $T_2$  are related by

$$\ln(T_2/T_1) = V\beta(P_2 - P_1)/C_P$$

You may assume that the volume of the block stays approximately constant during the compression.  
*Hint: Think about entropy  $S(P,T)$ . What does reversible and adiabatic mean for entropy? Then try to obtain an expression involving  $T$  and  $P$  (the variables mentioned in the question...)*

**5. From Gibbs function to equation of state**

A gas has molar Gibbs Free energy given by

$$g = RT \ln P + A + BP + \frac{1}{2}CP^2 + \frac{1}{3}DP^3$$

where  $A$ ,  $B$ ,  $C$  and  $D$  are constants. Find the equation of state (i.e. the relationship between  $P$ ,  $V$ , and  $T$ ) and explain why it is independent of  $A$ .

**6. A harmonic material**

Suppose a material has an equation of state per kg given by

$$p = A(v - b) + CT$$

Given that at  $P = 0$ ,  $T=300\text{K}$ :  $v_0 = 10^{-3}\text{m}^3/\text{kg}$ ;  $K_T = 10^{10}\text{Pa}$ ;  $\beta = 10^{-5}\text{K}^{-1}$ , determine the constants  $A$ ,  $b$ ,  $C$ ? Is this a gas or a condensed phase?

**7. Challenge Question: Deriving the ideal gas equation from experimental laws.**

Use *Joule's Law* (the internal energy of an ideal gas depends only on temperature) and *Boyle's Law* (at constant temperature, the product of pressure and volume for a fixed amount of an ideal gas is a constant), to derive the form of the equation of state of the ideal gas.

*Hint: start with the Central Equation, and employ one of the Maxwell relations. Then integrate...*

## 7 Questions: Expansion Processes

### 1. Free expansion of the van der Waals gas

Derive the expression for the Joule coefficient of the van der Waals equation of state.

$$\left(\frac{\partial T}{\partial V}\right)_U = -\frac{a}{C_V} \left(\frac{n}{V}\right)^2$$

*Hint: this requires use of the cyclical relation, the Central Equation, and a Maxwell relation...*

### 2. Joule-Kelvin coefficients

- (a) The Joule-Kelvin expansion is an isenthalpic process, in which the change in temperature is given: by

$$\Delta T_{JK} = \int_{P_1}^{P_2} \left(\frac{\partial T}{\partial P}\right)_H dP$$

Show that the Joule-Kelvin coefficient can be written in terms of the heat capacity and the coefficient of thermal expansion:

$$\mu_{JK} = \left(\frac{\partial T}{\partial P}\right)_H = \frac{V}{C_P}(\alpha T - 1)$$

- (b) Show that the Joule-Kelvin coefficient is zero for an ideal gas.  
 (c) Evaluate the Joule-Kelvin coefficient for a van der Waals gas with specific heat capacity  $c_P$ .  
 (d) Show that, for a van der Waals gas, the line dividing positive and negative values of  $\mu_{JK}$  (the “inversion curve”) is given by:

$$T_{in} = \frac{2a}{Rb} \left(\frac{v_{in} - b}{v_{in}}\right)^2$$

*Hint: The easiest way to get an expression for  $(\partial v/\partial T)_P$  from the van der Waals equation is to write it as  $T(P, V)$  and find  $(\partial T/\partial v)_P$ .*

#### (e) Cooling in an adiabatic expansion

Draw two isotherms and an adiabat for an ideal gas on an indicator ( $PV$ ) diagram, and hence show that an ideal gas always cools when undergoing a reversible adiabatic expansion. You do not need to do any detailed mathematics.

Now show that the coefficient for cooling in an adiabatic expansion is:

$$\left(\frac{\partial T}{\partial P}\right)_S = \frac{T}{C_P} \left(\frac{\partial V}{\partial T}\right)_P$$

Hence explain why an adiabatic expansion produces more cooling than an isenthalpic process, with a similar pressure change.

- (f) Discuss why the isenthalpic Joule-Kelvin process is used in real refrigerator, rather than the adiabatic or Joule processes.

### 3. Expanding through a phase transition

In a throttling process we normally write

$$\Delta T_{JK} = \int \left(\frac{\partial T}{\partial P}\right)_H dP = \int \frac{V}{C_P} (\alpha T - 1) dP$$

In an evaporator, the cooling substance goes from liquid to gas in an isenthalpic process. With reference to the equation above, explain why this is a good thing to do, and why it creates a problem applying the equation. What happens in practice is that the system remains at constant pressure and temperature, while extracting heat from the surroundings to overcome latent heat and allow it to evaporate.

**4. Gibbs Helmholtz**

A system is heated from temperature  $T_1$  to  $T_2$ , show that

$$G_2 = \frac{G_1 T_2}{T_1} - T_2 \int_{T_1}^{T_2} \frac{H}{T^2} dT$$

If the interatomic forces are known, it is fairly simple to compute the internal energy of a system. Suppose a material at atmospheric pressure in its liquid phase has internal energy

$$U = U_l(V) + A_l T$$

and in its solid phase

$$U = U_s(V) + A_s T$$

Assume further that it has zero thermal expansion:  $(\frac{dV}{dT})_P = 0$  and melting point of 300K.

Write down the gibbs free energy difference at 300K, and calculate its value at any other temperature  $T_2$ . Discuss each term in your final expression. By examining what happens at  $T = 0$ , prove that the given expressions for  $U$  are no longer valid in that limit.

5. **Critical point** At the critical point, where liquid and gas become indistinguishable,  $(\partial P / \partial V)_T = 0$  and  $(\partial^2 P / \partial V^2)_T = 0$ . Sketch isotherms on a PV diagram showing these two conditions. Show that, for a van der Waals gas, the critical point is at:

$$P_c = \frac{a}{27b^2}, \quad V_c = 3nb, \quad T_c = \frac{8a}{27Rb}$$

For a van der Waals gas for which  $a = 0.27 \text{ N m}^4 \text{ mole}^{-2}$  and  $b = 3.1 \times 10^{-5} \text{ m}^3 \text{ mole}^{-1}$  (approximately the values for carbon dioxide) calculate the values of  $P_c$ ,  $V_c$  and  $T_c$ . Compare the value of the critical temperature  $T_c$  with the value of the Boyle temperature  $T_B$  and comment.

## 8 Questions: Thermodynamics in other systems

### 1. A rubber band

- (a) Derive the so-called ‘energy equation’

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial P}{\partial T}\right)_V - P$$

for a simple substance. If the equation of state of the substance is known, the energy equation allows one to calculate the dependence of the internal energy on volume.

*Hint: Use the Maxwell relation derived from the Helmholtz function.*

- (b) Write down the analogous result for rubber band, where work done is tension times extension. Why is it different from force being the differential of energy?
- (c) The equation of state of a rubber band is

$$\mathcal{F} = aT \left( \frac{L}{L_0} - \left( \frac{L_0}{L} \right)^2 \right)$$

where  $a$  is a constant and  $L_0$  is the unstretched length. For this band, show that  $U$  is a function of  $T$  only.

- (d) If  $L_0 = 1$  metre and  $a = 1.3 \times 10^{-2} \text{ N K}^{-1}$ , calculate the work done on the band and the heat rejected when it is stretched isothermally and reversibly from 1 m to 2 m at  $T = 300 \text{ K}$ .

## 2. An adiabatic variation on the rubber band.

If the rubber band in the previous problem is stretched *adiabatically* and reversibly from 1 m to 2 m, by how much does its temperature rise? (Take the heat capacity to be  $C_L = 1.2 \text{ J K}^{-1}$ .)

## 3. Entropy of diamond

The low temperature specific heat of diamond varies with temperature as:

$$c_p = 124 \left( \frac{T}{\theta_D} \right)^3 \text{ kJ kg}^{-1} \text{ K}^{-1}$$

where the Debye temperature  $\theta_D = 1860 \text{ K}$ . What is the entropy change of 1 g of diamond when it is heated at constant pressure from 4 K to 300 K?

The Debye temperature of graphite is  $\theta_D = 1500 \text{ K}$ , and

$$c_p = 89 \left( \frac{T}{\theta_D} \right)^3 \text{ kJ kg}^{-1} \text{ K}^{-1}$$

Given that diamond has higher density than graphite, discuss whether the phase boundary on a PT phase diagram has positive or negative slope? Does the equation for specific heat satisfy the Third Law, that the change in entropy in any process becomes zero at 0K?

(The atomic weight of carbon is 12.)

## 4. Planck's Law for frequencies

Taking care about units and definitions, use the relation  $\nu = c/\lambda$ , rewrite the Planck distribution

$$u_\lambda(\lambda, T) = \frac{hc}{k_B \lambda^5} \left( \frac{1}{e^{hc/\lambda k_B T} - 1} \right)$$

in terms of frequency:

$$u_\nu(\nu, T) = \frac{2h\nu^3}{c^2} \frac{1}{\exp(h\nu/k_B T) - 1}$$

## 5. Planck's Law

The equation of state for radiation in a cavity is  $P = U/3V$ , where  $U(T)$  is the energy and  $V$  the volume. Use this with the Central Equation to show that the energy density varies as the fourth power of temperature.

Starting from Planck's ansatz relation between  $u$  and  $s$   $\frac{\partial^2 s}{\partial u^2} = \frac{c_1(\lambda)}{u(u+c_2(\lambda))}$

and integrating, derive the Planck distribution  $u_\lambda(\lambda, T)$ . You will need to use the Rayleigh and Wein limits to reintroduce  $\lambda$  via  $c_1$  and  $c_2$

The total energy density must be the integral of the energy density over all frequencies.

Show that the Planck distribution of energy between wavelengths satisfies the requirement that the total energy density varies as the fourth power of temperature. Show that the classical distribution ( $u_\lambda(\lambda, T) = k_B T \lambda^{-4}$ ) with each mode having  $\lambda$ -independent energy  $k_B T$  gives divergent energy.

What would the Boltzmann distribution give?

## 6. Photon Gas

- For a photon gas, the equation of state can be written  $P = \frac{u(T)}{3}$  where  $u(T)$  is the specific internal energy, which depends only on temperature. Show that the internal energy of a photon gas varies as the fourth power of temperature,  $U = kVT^4$ , where  $k$  is a constant.
- An evacuated cylinder of volume  $1\text{ m}^3$  contains a photon gas confined by a piston. At what temperature would the cylinder need to be for the piston to move outward against atmospheric pressure ( $10^5 \text{ Pa}$ ) ( $k = 7.56 \times 10^{-16} \text{ J/m}^3/\text{K}^4$ )

- (c) Show that the Heat Capacity  $C_v$  of a photon gas obeys the Third Law, and evaluate the entropy of a photon gas, assuming that  $S=0$  at  $T=0$ .
- (d) Show that the Gibbs Free Energy of a photon gas is zero and comment on the implication for creation and absorption of photons at a cavity wall.
- (e) Using the fact that  $G = 0$ , and *without* using the equation of state, prove that the pressure of a photon gas depends on temperature only.

### 7. Temperature of the sun.

By assuming that both the earth and the sun are perfect black bodies that radiate equally in all directions use the Stefan Boltzmann law to estimate the temperature of the sun. (You will need Temperature of the earth  $\approx 287K$ , Radius of the sun  $\approx 6.96 \times 10^8 m$ , Distance from earth to sun (astronomical unit)  $\approx 1.5 \times 10^{11} m$ ).

Comment on the various assumptions you have made.

*Hint: Think of how much area the earth takes up relative to the total emission surface of the sun at the orbital distance. Assume there is a radiative equilibrium on the earth. Note the earth's radius cancels out during this calculation.*

*Historical note: Stefan, lacking accurate knowledge of the relevant distances, performed this calculation by using a measure of the incident flux on earth. This was measured using huge reflecting mirrors and target objects!*

### 8. Black hole entropy

A black hole has only three independent variables, mass, charge and angular momentum. When a particle falls into a black hole, information (entropy) is “lost”.

It has been predicted that the entropy of a (non-rotating, uncharged) black hole depends on a single variable, such as its surface area (A) measured in Planck areas,  $S = k_B A c^3 / 4 G \hbar$ . This is called the “Bekenstein-Hawking formula”. Use it to calculate the temperature of a mini black hole of mass  $M = 10^{12}$  kg, and a black hole the size of the sun  $M = 2 \times 10^{30}$  kg.

*Hint: You will need to identify, and find values for,  $k_B$ ,  $G$ ,  $\hbar$  and  $c$ . The radius of a black hole depends only on its mass  $r = 2Gm/c^2$  (Schwartzchild radius), and its energy also depends only on the mass  $U = mc^2$ . The Central Equation of Thermodynamics lets you introduce  $T$  explicitly. The same equation suggests that the appropriate variables are  $U$  and  $V$ , so that **formally**  $S = S(U, V)$ . Assume that  $S$  is a function of  $U$  only.*

## 9 Questions: Phase transitions

### 1. Phases and Clausius-Clapeyron


- (a) Sketch the P-T projection of the PVT surface for a simple substance and identify its main features.
- (b) By using the fact that the specific Gibbs functions of coexisting phases are the same, derive the Clausius-Clapeyron equation

$$\frac{dp}{dT} = \frac{l}{T(\nu_2 - \nu_1)}$$

where  $\nu_1$  and  $\nu_2$  are the specific volumes of the two phases and  $l$  is the specific latent heat of the transition.

- (c) In a pressure cooker, food is cooked more quickly by raising the pressure so that water boils at a higher temperature. Estimate the pressure at which water boils at  $130^\circ\text{C}$ . The density of water and steam at  $100^\circ\text{C}$  are  $959 \text{ kg m}^{-3}$  and  $0.600 \text{ kg m}^{-3}$  respectively, and the latent heat of vaporisation is  $2257 \text{ kJ kg}^{-1}$ . Take 1 atmosphere to be  $10^5 \text{ N m}^{-2}$ .

## 2. Phase mixtures



The figure shows an isotherm on the  $P-V$  projection. At the point X the substance is a mixture of liquid and vapour. The total mass of substance is the sum of the liquid and vapour masses  $m = m_l + m_v$ . At coexistence  $v_l$  and  $v_v$  are the specific volumes of the liquid and vapour (per kg).

Write down an expression for the total volume at X.

Let the specific volume of the mixture at X be  $v$ . Show that :

$$m_l(v - v_l) = m_v(v_v - v)$$

Explain why this result, which gives the ratio  $m_v/m_l$ , is known as the 'lever rule'.

How does the Gibbs free energy change with volume in the coexistence region?

What is the compressibility of the substance at X.

## 3. Solid-solid phase transitions

Tin can exist in two forms ("allotropes"): Grey tin is the stable form at low temperatures and pressures. There is a first-order transition between the two phases at 291 K and 1 atm. Given the latent heat for the transition ( $18.5 \times 10^3 \text{ J kg}^{-1}$ ); and the densities  $5.75 \times 10^3 \text{ kg m}^{-3}$  (grey) and  $7.30 \times 10^3 \text{ kg m}^{-3}$  (white); estimate the change in this transition temperature if the pressure is increased to 100 atm?

*Hint: The grey tin to white tin transition is a phase transition, so use the Clausius Clapeyron equation. Try assuming the phase boundary is a straight line, putting  $dP/dT = \Delta P/\Delta T$  even though  $\Delta P = 99 \text{ atm}$ . Note also that the density change going from the low-temperature-stable grey tin to the high-temperature-stable white tin is an increase, like in ice-to-water – so anticipate  $dP/dT$  to be negative.*

## 4. Triple Point

The sublimation and the vaporization curves of a particular material are given by:

$$\ln P = 0.04 - 6/T \quad \text{sublimation}$$

$$\ln P = 0.03 - 4/T \quad \text{vaporization}$$

where  $P$  is in atmospheres.

- Find the temperature and pressure of the triple point.
- By assuming that the volume in the gas phase is much larger than for the solid or liquid, show that the latent heats are  $4R$  and  $6R$  per mole.
- By considering entropy around a cycle around the triple point in the  $PT$  phase diagram, show that

$$\frac{l_{SV}}{T_{TP}} - \frac{l_{SL}}{T_{TP}} - \frac{l_{LV}}{T_{TP}} = 0$$

and hence calculate the latent heat of fusion.



*Hint: Consider a loop round the triple point in the  $PT$  phase diagram. As  $S$  is a state function,*

$$\frac{l_{SV}}{T_{TP}} - \frac{l_{SL}}{T_{TP}} - \frac{l_{LV}}{T_{TP}} = 0$$

## 5. Impossible Phase diagram

Consider the phase diagram shown below on the left, in which three proposed phases are separated by first order phase transitions. A diamond anvil cell experiment applies an isothermal pressure increase which traverses the three phases.

- List the phases 1-3 in order of density. Comment on the compressibilities.
- By considering isothermal pressure increase just below the triple point, show that this phase diagram is impossible.
- The diagram on the right represents a shock compression process, along the line AB, which has constant slope  $m$ . Define the quantity (call it  $Z$ ) which is conserved along this line. Is this quantity a state variable?
- By generalising parts (a) and (b) to consider  $Z$  and its conjugate variable, show that no sector of the phase diagram can have an angle greater than  $180$  degrees at the triple point. You do not need evaluate the conjugate variable.



#### 6. Pressure effect on melting ice

Water has the unusual property that the solid phase (ice) can be melted by applying sufficiently large pressure. Use the Clausius-Clapeyron equation to determine the additional pressure above atmospheric  $\Delta P$  required to melt ice at temperature  $-\Delta T$  below freezing (at  $0^\circ\text{C}$ ). This value is in effect the slope of the phase boundary in the  $P$ - $T$  representation. Comment on the sign of the value. Take  $l$  the latent heat of fusion to be  $333.7\text{kJkg}^{-1}$ , and the densities of water and ice at  $0^\circ\text{C}$  to be  $\rho_w = 1000\text{kgm}^{-3}$  and  $\rho_i = 916\text{kgm}^{-3}$ .

Assuming the slope of the phase boundary is constant away from the atmospheric melting point of water (which turns out to be a valid assumption), use your calculated value in the following situations.

- Melting glaciers. The bottom surface of a  $2000\text{m}$  thick glacier is at a temperature of  $-1.5^\circ\text{C}$ . Will the water here be liquid or ice? Use the density of ice given above.
- Very cold water. The lowest temperature at which liquid water is stable is  $-22^\circ\text{C}$ . What pressure must be applied to keep the water from freezing? Express your answer in Pa and atm.
- Ice skates. Do ice skates apply sufficient pressure on the surface of ice to melt it? You will have to make some assumptions regarding the area of skate in contact with the ice, the temperature and the weight of the skater

## 10 Questions: Chemical Potential

### 1. Irn Bru

A 1.05l bottle of Irn Bru contains 1l of drink and is pressurised to 5atm with CO<sub>2</sub> gas.

At this pressure, the CO<sub>2</sub> concentration in the drink is given by Henry's law  $C = 0.031 P_g$  where  $C$  is the concentration in moles/L, and  $P_g$  is the partial pressure of the gas (5.0 atm).

What is the chemical potential of the CO<sub>2</sub> in this system relative to the gas at atmospheric pressure?

The cap is briefly loosened, so that the gas comes fully into equilibrium with the atmosphere, but the dissolved CO<sub>2</sub> remains in solution. The bottle is then sealed.

What are the chemical potentials of the CO<sub>2</sub> in the drink and the gas above?

Now the bottle is shaken, such that the contents come into equilibrium. Describe what happens, and calculate the pressure inside the bottle now?

Approximately how many times can this process be repeated before the drink goes flat?

### 2. Chemical Potential: Nature's boundary condition

Show that the molar chemical potential for an ideal gas at temperature  $T$  is given by

$$\mu = RT \ln(P/P_0) + \mu_0$$

where  $\mu_0 = \mu(T, P_0)$ .

Under standard temperature and pressure, dilute carbon dioxide has  $\mu_0 = 394\text{kJ/mol}$  in air and  $\mu_0 = 386\text{kJ/mol}$  in water.

If the atmosphere contains 0.04% CO<sub>2</sub>, estimate the concentration of CO<sub>2</sub> dissolved in the ocean.

### 3. Chemical Potential Change in Mixing

A rigid, thermally isolated container holding 1mol of argon at 1atm, 300K is connected to an identical container holding 1mol of krypton at 1atm, 300K. No heat is added and no work is done on the system. Without calculation, explain what you expect to happen and the final equilibrium state. How do the pressure, temperature and entropy change?

How would the result change if the initial amounts were different, in volumes  $V_A$  and  $V_K$ , still at the same initial T and P and still totalling 2 moles of atoms?

### 4. Regular solution and solubility limits

A total of one mole of fluids, comprising two atomic types, A and B, are mixed at constant temperature. The fluids are ideal except that they repel one another, which adds a term to the internal energy  $\frac{Z}{v_A v_B}$ .

Explain why this is a reasonable form for the interaction.

Calculate the change in Gibbs Free Energy when the two are mixed, with mole fractions  $x_A$  and  $x_B = 1 - x_A$ , and plot this as a function of  $x_A$  for various values of  $\frac{RT}{Z}$ .

For  $Z = 3RT$  what values of  $x_A$  minimise  $\Delta g$ ? Calculate the chemical potential for species A at these values.

Describe the mixing process as  $x_A$  increases from 0 to 1.

### 5. Supercool

Salt is added to a mixture of ice and water at 0°C. Assuming that salt cannot dissolve in ice, what is the change in chemical potential of the water with salt concentration  $X = 0.1$ ?

$$\left(\frac{\partial \mu_L}{\partial X}\right)_T$$

What is the final temperature of the mixture? Assume that the partial pressure of each component in the brine is proportional to its concentration. Take the latent heat of fusion to be 18kJ/mol. For convenience, you can assume that for small changes in absolute temperature  $1/T^2$  is temperature independent.

### 6. Simplified Osmosis

A surface-dwelling single-celled, spherical marine creature contains protein molecules and water, and is separated from the sea by a semipermeable membrane<sup>1</sup> which water can pass, but not protein. Treating the protein as a monotonic ideal gas, and assuming there are 2% as many protein molecules as water molecules, calculate the total pressure inside the cell.

If the membrane has diameter 10 $\mu$ m and thickness 10nm, then what is the stress in the membrane?

---

<sup>1</sup>i.e. the cell wall. Real cells have all sorts of funny shaped or charged holes to let through some molecules and not others