

STATISTICAL PHYSICS 06/07

The Ising Model

Tutorial Sheet 6

The questions that follow on this and succeeding sheets are an integral part of this course. The code beside each question has the following significance:

- **K**: key question – explores core material
- **R**: review question – an invitation to consolidate
- **C**: challenge question – going beyond the basic framework of the course
- **S**: standard question – general fitness training!

6.1 **Mean-Field Theory of a Three State Model [S]** A set of spins on a lattice of coordination number z can take values $S_i = (1, 0, -1)$, as opposed to just $(1, -1)$ as in the Ising model. The configurational energy is (as usual)

$$E(\{S_i\}) = -J \sum_{\langle ij \rangle} S_i S_j - h \sum_i S_i$$

Show that in the mean-field approximation, the magnetization per site obeys

$$m = \frac{2 \sinh(\beta(Jzm + h))}{2 \cosh(\beta(Jzm + h)) + 1}$$

and find the critical temperature, T_c .

6.2 **Critical Exponents from Mean-Field Theory of Ising [R/S]**

(i) If we denote the order parameter (or magnetisation per spin) by m , show that the mean-field solution of the Ising model can be written as

$$m = \tanh \left[\frac{m}{1-t} + b \right],$$

where $t = (T_c - T)/T_c$ is the reduced temperature and $b = h/kT$ is the reduced external magnetic field. By considering m for temperature close to T_c and for zero external field, show that the associated critical exponent takes the value $\beta = \frac{1}{2}$.

[Hint: the following expansion

$$\tanh x = x - \frac{1}{3}x^3 + O(x^5),$$

for small values of x , should be helpful.]

(ii) Obtain an expression for the mean energy \bar{E} of the Ising model when the applied field is zero, using the simplest mean-field approximation. Hence show that the heat capacity C_h has the behaviour:

$$\begin{aligned} C_h &= 0, & \text{for } T > T_c; \\ &= 3Nk/2, & \text{for } T < T_c. \end{aligned}$$

Comment on the nature of the singularity at T_c . Is there a critical exponent α ?

- (iii) By considering the behaviour of the order parameter at temperatures just above the critical point, show that the critical exponent γ , which is associated with the isothermal susceptibility, takes the value $\gamma = -1$ according to mean-field theory.

Also, by considering the effect of an externally applied magnetic field at $T = T_c$, show that the exponent associated with the critical isotherm takes the value $\delta = 3$.

- 6.3 **The Lattice Gas [S]** Check the claim, made in the notes, that the Energy of the lattice gas model in the grand canonical ensemble

$$E'(\{c\}) = \epsilon \sum_{\langle ij \rangle} c_i c_j - \mu \sum_i c_i$$

is transformed to that of the Ising model:

$$E'(\{s\}) = -J \sum_{\langle ij \rangle} S_i S_j - h \sum_i S_i + \frac{\epsilon z - 4\mu}{8} \mathcal{N}$$

where \mathcal{N} is the number of sites, by means of the change of variable

$$S_i = 2c_i - 1 = \pm 1 \quad ; \quad -J = \frac{\epsilon}{4} \quad ; \quad -h = \frac{\epsilon z - 2\mu}{4}$$

- 6.4 **Transfer Matrix Solution of the one-dimensional Ising Model [C]**

Show that the canonical partition function of the one-dimensional Ising model (with periodic boundary conditions) may be written as

$$Z_c = \sum_{S_1=\pm 1} \sum_{S_2=\pm 1} \dots \sum_{S_N=\pm 1} T(S_1, S_2) T(S_2, S_3) \dots T(S_N, S_1)$$

where

$$T(S_i, S_{i+1}) = \exp \left[\beta J S_i S_{i+1} + \frac{\beta h}{2} (S_i + S_{i+1}) \right]$$

Hence show that

$$Z_c = \lambda_+^N + \lambda_-^N \simeq \lambda_+^N$$

where λ_{\pm} are the eigenvalues, which you should determine, of the transfer matrix \hat{T}

$$\hat{T} = \begin{pmatrix} T(1, 1) & T(1, -1) \\ T(-1, 1) & T(-1, -1) \end{pmatrix}$$

Show that the magnetisation is zero for $h = 0$ at all temperatures T and find an expression for the mean energy at low temperature.