

# Physical Mathematics 2010: Problems 2 (week 4)

## Curvilinear coordinate systems

1. Find the scale factors for changing volume integrals from Cartesian coordinates  $(x, y, z)$  to the following coordinate sets:

- (a) Cylindrical polar coordinates  $(\rho, \phi, z)$ :  $x = \rho \cos \phi$ ;  $y = \rho \sin \phi$ ;  $z = z$ .
- (b) Spherical polar coordinates  $(r, \theta, \phi)$ :  $x = r \sin \theta \cos \phi$ ;  $y = r \sin \theta \sin \phi$ ;  $z = r \cos \theta$ .

### 2. Cylindrical polars

- (a) Determine scale factors for cylindrical polar coordinates and the gradient operator in the  $\mathbf{e}_r, \mathbf{e}_\phi, \mathbf{e}_z$  basis.
- (b) Combine this with the divergence formula:

$$\nabla \cdot \mathbf{v} = \frac{1}{h_1 h_2 h_3} \left[ \frac{\partial h_2 h_3 v_1}{\partial \xi_1} + \frac{\partial h_3 h_1 v_2}{\partial \xi_2} + \frac{\partial h_1 h_2 v_3}{\partial \xi_3} \right]$$

to find the Laplacian in cylindrical polar coordinates.

- (c) Seek separable solutions in the cylindrical coordinates:

$$f(x, y, z) = R(r)\Phi(\phi)Z(z)$$

and separate the equation into three ODE's and separation constants.

- (d) Suppose  $f$  has boundary conditions in the  $z$ -direction such that

$$f(x, y, 0) = f(x, y, L) = 0$$

and that

$$\Phi(2\pi) = \Phi(0).$$

What constraints does this introduce on the allowed separation constants?

## Wave equation for light

3. If  $\nabla \cdot \mathbf{E} = 0$ ,  $\nabla \cdot \mathbf{B} = 0$ ,  $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$ , and  $\nabla \times \mathbf{B} = \frac{\partial \mathbf{E}}{\partial t}$  use the "GDMCC" rule for the vector Laplacian show that  $\mathbf{E}$  and  $\mathbf{B}$  satisfy a wave equation.

## Method of Froebenius & Special functions

### 4. Bill and Ted's excellent misadventure

Bill and Ted have brought Pythagoras to the future and lost him in a night club. We now live in a world without sin and cos.

To rectify this it is up to you to use the method of Froebenius to rediscover these precious functions

- (a) Substitute the infinite series  $y(x) = \sum_{n=0}^{\infty} C_n x^n$  to the differential equation

$$y'' + y = 0$$

You should end up with two sums.

- (b) Relabel the summation using  $m = n - 2$  on the  $y''$  term
- (c) Use a notation where  $C_i = 0$  for  $i < 0$  to sum the  $y$  term over the range  $\sum_{n=-2}^{\infty}$
- (d) Hence obtain the *indicial equation*

$$C_{m+2}(m+1)(m+2) = -C_m$$

This relates every other coefficient in a *recurrence relation*.

- (e) Deduce that  $C_0$  can be non-zero even though  $C_{-2} = 0$ , and that  $C_1$  can be non-zero even though  $C_{-1} = 0$ .  
We have two independent series, and so two free parameters  $C_0$  and  $C_1$  as should be the case for a 2nd order ODE.
- (f) Find the series with (a)  $C_0 = 1, C_1 = 0$  and, (b)  $C_0 = 0, C_1 = 1$
- (g) How would you make up the world's first table of sinusoids?
- (h) Give these two independent series their names to save the world from Bill and Ted's misadventure.
5. **Exponential:** apply the above method to find the exponential function solving the 1st order ODE

$$y' = y$$

fixing the single free parameter via  $y(0) = 1$

6. **Bessel bookwork:** use the method in the notes to solve Bessel's equation in the case  $n = 0$ , and  $k = 1$ :

$$r^2 R'' + rR' + r^2 R = 0$$

Solve for the first four terms of the series solution that remains finite at  $r = 0$ , and sketch the behaviour of the function.

7. **Cauchy's Equation:**

$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{n(n+1)}{r^2} R = 0$$

Verify (by substitution) that it is solved by solutions of the form  $R(r) = r^p$  for  $p = n$  or  $p = -(n+1)$ . Explain why the general solution is simply a linear combination of these. [i.e. Why is it a linear combination? Why are there no more functions?]

8. By making the substitutions  $x = pr$ , show that the following radial equation for a function  $R(r)$ :

$$r^2 R'' + rR' + (p^2 r^2 - n^2) R = 0$$

can be written in the standard form of Bessel's Equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2)y(x) = 0. \quad (1)$$

The solutions to Bessel's equation for give (fixed)  $n$  are  $y(x) = J_n(x)$ , which are the Bessel functions of order  $n$ .

By substituting into Bessel's equation, show the following are Bessel functions and find their order:

$$(i) \quad \frac{\sin x}{\sqrt{x}} \quad (ii) \quad \frac{\cos x}{\sqrt{x}}$$

Explain why these solutions will never be seen in when solving the radial equation in problems with circular or cylindrical symmetry.

9. A circular membrane has its edge attached to a fixed circular ring centred on the origin and of radius unity. For small displacements from its equilibrium position  $u(\rho, \phi, t)$  perpendicular to the plane of the ring

$$\frac{1}{\rho} \frac{\partial}{\partial \rho} \left( \rho \frac{\partial u}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2 u}{\partial \phi^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} ,$$

for times  $t > 0$  where  $0 \leq \rho \leq 1$  and  $0 \leq \phi < 2\pi$  are radial and angular plane-polar coordinates respectively.

If the motion is axially symmetric, explain why we can solve this differential equation by considering solutions of separated form  $u(\rho, \phi, t) = R(\rho) T(t)$ .

Show that the radial equation reduces to

$$\frac{d^2 R}{d\rho^2} + \frac{1}{\rho} \frac{dR}{d\rho} + k^2 R = 0$$

where  $k$  is a constant.

By making the substitution  $x = k\rho$ , explain why the solution is  $R(\rho) = J_0(k\rho)$ , given that  $J_0(x)$  is the only solution of the differential equation

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0$$

that is finite at the origin.

Apply the boundary conditions to show that the general solution is

$$u(\rho, \phi, t) = \sum_{n=1}^{\infty} \{ E_n J_0(\lambda_n \rho) \cos(\lambda_n ct) + F_n J_0(\lambda_n \rho) \sin(\lambda_n ct) \} ,$$

where  $E_n$  and  $F_n$  are unknown constants and the zeros of  $J_0$  are located at  $\lambda_n$  for  $n = 1, 2, \dots$ , i.e.  $J_0(\lambda_n) = 0$ .

Initially the membrane is held at rest in the shape of a cone with its centre distance  $h$  above its equilibrium position. If the membrane is then released, show that the displacement of the point  $(\rho, \phi)$  of the membrane at a later time is given by

$$u(\rho, \phi, t) = \sum_{n=1}^{\infty} A_n \cos(\lambda_n ct) J_0(\lambda_n \rho) ,$$

where

$$A_n = \frac{2h}{[J_0'(\lambda_n)]^2} \int_0^1 d\rho \rho(1-\rho) J_0(\lambda_n \rho) .$$

[The relation

$$\int_0^1 ds s J_0(\lambda_n s) J_0(\lambda_m s) = \frac{1}{2} \delta_{mn} [J_0'(\lambda_n)]^2$$

may be assumed without proof, where  $\delta_{mn} = 1$  for  $m = n$  and zero otherwise.]

(*Maths Skills, Dec.08*)

## 10. Quantum square well

The Schrodinger equation in a two dimensional infinite square well defined by

$$V(x, y) = \begin{cases} 0 & : x, y \in [0, L] \\ \infty & \text{otherwise} \end{cases}$$

is

$$i\hbar \frac{\partial}{\partial t} \psi(x, y, t) = -\frac{\hbar^2}{2m} \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \psi(x, y, t) + V(x, y) \psi(x, y, t).$$

(a) Substitute a separable solution of the form  $X(x)Y(y)T(t)$  and derive the separate ODE's for  $X(x)$ ,  $Y(y)$ , and  $T(t)$ .

(b) Apply the boundary conditions

$$\psi(x, L, t) = \psi(L, y, t) = \psi(x, 0, t) = \psi(0, y, t) = 0$$

to  $x$  and  $y$  and find the allowed values for the separation constants governing the  $X$ ,  $Y$  and  $T$  differential equations.

(c) Suppose

$$\psi(x, y, t = 0) = x(L - x)y(L - y).$$

Find the wavefunction  $\psi$  for all later times.

*Maths Skills Nov 2010*

## 11. Drumskin

a) The displacement  $u(x, y, t)$  of a circular drumskin of radius  $L$  is described in Cartesian coordinates by the two dimensional wave equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2}.$$

In circular polar coordinates  $(r, \theta)$  the Laplacian is

$$\nabla^2 f = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial f}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 f}{\partial \theta^2}$$

Rewrite the wave equation in circular coordinates, and use the method of separation of variables to determine separated differential equations in  $r$ ,  $\theta$ , and  $t$  and define the separation constants.

b) Solve the  $\theta$  equation and apply boundary conditions to show that the radial equation reduces to Bessel's equation of integral order

$$r^2 R'' + rR' + (k^2 r^2 - n^2)R = 0$$

Where  $R(r)$  is the separated radial dependence,  $k$  is a separation constant, and  $n$  is an integer.

c) The solutions are Bessel functions of the form  $R(r) = J_n(kr)$ .

Apply boundary conditions in the  $r$  variable to determine the frequency of the corresponding normal mode in terms of  $\alpha_{nm}$ , where  $\alpha_{nm}$ , is the  $m$ -th zero of the Bessel function  $J_n$ .

d) The zeroes of the first two Bessel functions are

$$\begin{array}{cc} J_0 & J_1 \\ \alpha_{01} = 2.405 & \alpha_{11} = 3.832 \\ \alpha_{02} = 5.520 & \alpha_{12} = 7.016 \end{array} .$$

A drummer strikes the drum somewhere with  $\theta = 0$ . Considering only these four modes, where should he hold his other drumstick to excite:

i) only the  $n = 0, m = 2$  mode?

ii) both the  $n = 1$  modes with no  $n = 0$  contributions?

iii) only the  $n = 1, m = 2$  mode?

*Physical Mathematics May 2011*

## 12. Hermite Polynomials & and Harmonic Oscillator

The time independent Schrödinger equation for a particle of mass  $m$  in a harmonic oscillator potential  $V(x) = \frac{1}{2}kx^2$  can be placed in the form

$$\frac{d^2}{dx^2}\psi(x) + (\lambda - x^2)\psi(x) = 0,$$

where  $\lambda = \frac{2E}{\hbar\omega}$ ,  $E$  is the energy, and the angular frequency  $\omega = \sqrt{\frac{k}{m}}$ .

- (a) Show that after a substitution  $\psi(x) = e^{-\frac{x^2}{2}}H(x)$  the equation reduces to

$$H'' - 2xH' + (\lambda - 1)H = 0.$$

- (b) Substitute a power series for  $H(x)$ :

$$H(x) = \sum_{n=-\infty}^{\infty} c_n x^n.$$

Use the method of Fröbenius to determine a recurrence relation for  $c_n$ .

- (c) Analyse this recurrence relation:

For what values of  $n$  can the series begin and terminate?

What is the form of  $\psi(x)$  at large  $x$ , and how does normalisability constrain  $\lambda$ ?

- (d) Deduce (unnormalised) eigenfunctions  $\psi_1(x)$ ,  $\psi_2(x)$   $\psi_3(x)$  and associated energies  $E_1$ ,  $E_2$ ,  $E_3$  for the first, second and third states.

- (e) Consider operators  $a_+ = (x - \frac{d}{dx})$ , and  $a_- = (x + \frac{d}{dx})$ .

Apply  $a_+$  and  $a_-$  to your solution  $\psi_1$ .

Show that if  $\psi_n$  is a solution of the equation for  $\lambda = \lambda_n$ , then  $\psi_{n+1} = a_+\psi_n$  is a solution for  $\lambda = \lambda_n + 2$ .