


# Physics 3: Particle Physics

## Lecture 7: Introduction to the Weak Force March 3rd 2008



- \*Weak interactions
- \*Charged and neutral current
- \*Feynman Rules for weak force

1

## Introduction to the Weak Force

The weak force is responsible for some of the most important phenomena:

- Decays of the muon and tau leptons
- Neutrino interactions
- Decays of the lightest mesons and baryons
- Radioactivity, nuclear fission and fusion

Characteristics of Weak Processes:


- Long lifetimes  $10^{-13} - 10^3$  s
- Small cross sections  $10^{-13}$  mb

Boson	$W^\pm$	$Z^0$
Mass $\text{GeV}/c^2$	80.4	91.2
charge, $e$	$\pm 1$	0
spin	$1\hbar$	$1\hbar$

Weak Force is propagated by massive  $W^+$ ,  $W^-$  and  $Z^0$  bosons



- The interactions of  $W^\pm$  and  $Z^0$  are different (related by symmetry of the weak interaction)

- $W^\pm$  and  $Z^0$  can interact with each other
- $W^\pm$  and  $\gamma$  can interact (as  $W^\pm$  bosons are charged)



2

# Weak Vertices


QED	$W$ -boson
mediated by the exchange of virtual photons	mediated by the exchange of $W$ boson
acts on all charged particles	acts on all quark and leptons
coupling strength $\propto e \propto \sqrt{\alpha}$	coupling strength $\propto g_W \propto \sqrt{\alpha_W}$
propagator term: $1/(q^2 - m_\gamma^2) = 1/q^2$	propagator term: $1/(q^2 - m_W^2)$
For many processes: $\mathcal{M} \propto e^2/q^2$	For many processes: $\mathcal{M} \propto g_W^2/(q^2 - m_W^2)$
	

Recall: matrix element,  $\mathcal{M}$ , is the amplitude of a process.  
Scattering cross section,  $\sigma \propto \mathcal{M}^2$ . Decay width,  $\Gamma \propto \mathcal{M}^2$

3

# Interactions of the $W^\pm$ boson

- Known as “charged current interactions”
- Charged current acts on **all** fermions - quarks and leptons
- Charged current **changes the flavour of the fermion**:
  - e.g. electron emitting an  $W$ -boson can't remain an electron - violates conservation of charge!
    - an electron turns into a electron neutrino
    - an up quark turns into a down quark and vice versa!
- Coupling strength at every vertex  $\propto g_W$
- Propagator term describing the  $W$ -boson  $\propto \frac{1}{(\underline{q}^2 - m_W^2)}$ 
  - $\underline{q}$  is the four-momentum transferred by the  $W$ -boson



4

# Allowed Flavour Changes

At a  $W$ -boson vertex:

- Lepton numbers:  $L_e, L_\mu$  and  $L_\tau$ , is conserved:

Allowed lepton flavour changes:  $e^- \leftrightarrow \nu_e$   $\mu^- \leftrightarrow \nu_\mu$   $\tau^- \leftrightarrow \nu_\tau$

- Baryon number,  $B$ , is conserved

- Strangeness, Charmness ...  $S, C, B, T$  are violated:


Allowed quark flavour changes:

$(Q=+2/3 \text{ } e \text{ quark}) \leftrightarrow (Q=-1/3 \text{ } e \text{ quark})$

$(d \ s \ b) \leftrightarrow (u \ c \ t)$

- Each of the nine possible quark flavour changes has a different coupling strength, given matrix term  $V_{CKM}$
- Main quark flavour changes are within generations:

$d \leftrightarrow u$   $s \leftrightarrow c$   $b \leftrightarrow t$




5

# Interactions of the $Z^0$ boson

- Known as “neutral current interactions”
- Acts on **all** fermions - quarks and leptons
- Neutral current **conserves** flavour of the fermion
- No allowed fermion flavour changes

- Propagator term  $\propto \frac{1}{(q^2 - m_Z^2)}$

- Coupling depends on fermion flavour - we won't consider this in this course



Anywhere a photon could be exchanged a  $Z^0$  boson can be exchanged.  
(Almost vice-versa, except  $Z^0$  boson also has neutrino interactions too!)

**Electromagnetic and weak neutral current interactions are linked!**


6

# Feynman Rules for Weak Interaction

How to calculate the matrix element,  $\mathcal{M}$ , for a weak decay or scattering

e.g. decay of a muon  $\mu^- \rightarrow e^- \nu_\mu \bar{\nu}_e$

- Draw the Feynman diagram for the process
- give a four momentum for each particle



- Check quantum numbers conservation at every vertex
  - For both  $W$  and  $Z$ :  $L_e, L_\mu$  and  $L_\tau, B, Q$
  - For  $Z$  only: no change of quark or lepton flavour
- Is energy and momentum conserved? For decay:  $\sum m_{\text{initial}} > \sum m_{\text{final}}$
- Write down the coupling at each vertex:  $g_W$  (for  $W$ )
- Work out four-momentum transferred by boson:  $\underline{q} = (\underline{p}_3 - \underline{p}_1) = (\underline{p}_4 + \underline{p}_2)$
- Write down the propagator term for each boson:  $1/(\underline{q}^2 - m_{\text{boson}}^2)$
- $\mathcal{M}$  is proportional to product of vertex and propagator terms:  $\mathcal{M} \propto \frac{g_w^2}{(\underline{q}^2 - m_W^2)}$

7

## Summary

<p>The weak force acts on <b>all</b> quarks and leptons.</p> <p>Two massive bosons propagate the weak interaction: <math>W^\pm</math> and <math>Z^0</math>.</p>	<p>Weak interactions are characterised by:</p> <ul style="list-style-type: none"> <li>• Long lifetimes <math>10^{-13} - 10^3</math> s</li> <li>• Small cross sections <math>10^{-13}</math> mb</li> </ul>
<p><math>W^\pm</math>-boson interactions changes fermion flavour</p> <p><math>e^- \leftrightarrow \nu_e \quad \mu^- \leftrightarrow \nu_\mu \quad \tau^- \leftrightarrow \nu_\tau</math>  <math>(Q=+2/3 \text{ e quark}) \leftrightarrow (Q=-1/3 \text{ e quark})</math></p> <ul style="list-style-type: none"> <li>• quark coupling at <math>W^\pm</math> vertex: <math>g_W V_{CKM}</math></li> <li>• lepton coupling at <math>W^\pm</math> vertex: <math>g_W</math></li> <li>• <math>W^\pm</math> propagator term: <math>\frac{1}{(\underline{q}^2 - m_W^2)}</math></li> </ul>	<p><math>Z^0</math>-boson interactions conserve the flavour of the fermion</p> <ul style="list-style-type: none"> <li>• <math>Z^0</math>-boson propagator term: <math>\frac{1}{(\underline{q}^2 - m_Z^2)}</math></li> </ul> <p><math>Z^0</math>-boson interaction is connected to electromagnetic interaction</p>

8